

SHORT TITLE: GmYSL7, a peptide transporter important for SNF

Corresponding author details: Penelope Smith, School of Life Sciences, Agribio, 5 Ring Rd, Bundoora, Victoria, 3083, Australia.

TITLE: Soybean Yellow Stripe-like 7 is a symbiosome membrane peptide transporter important for nitrogen fixation

Alexander Gavrín², Patrick C. Loughlin², Ella Brear², Oliver W. Griffith⁶, Frank Bedon¹, Marianne Suter Grottemeyer⁴, Viviana Escudero⁵, Maria Reguera⁵, Yihan Qu², Siti N Mohd-Noor², Chi Chen², Marina Borges Osorio¹, Doris Rentsch⁴, Manuel González-Guerrero⁵, David A. Day³, Penelope Mary Collina Smith^{1*}.

¹School of Life Sciences. La Trobe University, Bundoora, Victoria 3083. Australia.

²School of Life and Environmental Science, The University of Sydney, Sydney, New South Wales, 2006, Australia

³College of Science and Engineering, Flinders University, Bedford Park, Adelaide, SA, Australia

⁴IPS, Molecular Plant Physiology, University of Bern, Altenbergrain 21, 3013 Bern Switzerland

⁵Centro de Biotecnología y Genómica de Plantas (UPM-INIA). Universidad Politécnica de Madrid. Campus de Montegancedo. Crta. M-40 km 38. 28223 Pozuelo de Alarcón (Madrid), Spain

⁶Department of Biological Sciences, Macquarie University, Macquarie Park, NSW, 2109

One sentence summary: Soybean Yellow Stripe-like 7 encodes an oligopeptide transporter that localises to the symbiosome membrane in nodules and contributes to nitrogen fixation and symbiosome development.

Author contributions: AG completed the gene silencing, RNAseq analysis and contributed to the localisation. PCL cloned the gene and contributed to the

localisation, promoter GUS analysis and real time analysis. EMB, SNMN, MSG, DR and PMCS completed the analysis of YSL7 in yeast. OG, EMB, MBO, FB and PMCS analysed the RNAseq results. VE, MR and MGG completed the complementation of MtYSL7. YQ completed the real time analysis. CC completed the promoter GUS analysis. PMCS, FB, EMB and DAD isolated symbiosome and microsomal membrane and did proteomic analysis. PMCS and DAD conceived the project and were involved in experimental design and analysis. PMCS and AG wrote the manuscript. All authors contributed to editing of the manuscript.

Funding information : This research was funded by the Australian Research Council Discovery Projects DP0772452, DP120102780 and DP150102264 and Industrial Transformation Research HUB IH140100013.

ABSTRACT

Legumes form a symbiosis with rhizobia that convert atmospheric nitrogen (N_2) to ammonia and provide it to the plant in return for a carbon and nutrient supply. Nodules, developed as part of the symbiosis, harbor rhizobia that are enclosed in a plant-derived symbiosome membrane (SM) to form an organelle-like structure called the symbiosome. In mature nodules exchanges between the symbionts occur across the SM. Here we characterize Yellow Stripe-like 7 (GmYSL7), a Yellow stripe-like family member localized on the SM in soybean (*Glycine max*) nodules. It is expressed specifically in infected cells with expression peaking soon after nitrogenase becomes active. Unlike most YSL family members, GmYSL7 does not transport metals complexed with phytosiderophores. Rather, it transports oligopeptides of between four and 12 amino acids. Silencing *GmYSL7* reduces nitrogenase activity and blocks infected cell development so that symbiosomes contain only a single bacteroid. This indicates the substrate of YSL7 is required for proper nodule development, either by promoting symbiosome development directly or by preventing inhibition of development by the plant. RNAseq of nodules where *GmYSL7* was silenced suggests that the plant initiates a defense response against rhizobia with genes encoding proteins involved in amino acid export down-regulated and some transcripts associated with metal homeostasis altered. These changes may result from the decrease in nitrogen fixation upon *GmYSL7* silencing and suggest that the peptide(s) transported by *GmYSL7* monitor the functional state of the bacteroids and

regulate nodule metabolism and transport processes accordingly. Further work to identify the physiological substrate for GmYSL7 will allow clarification of this role.

INTRODUCTION

Legumes form a symbiosis with soil bacteria, rhizobia, that allows them to access N₂ from the atmosphere. This symbiosis is an important contributor to the biological nitrogen cycle. The rhizobia fix N₂ via the enzyme nitrogenase to produce ammonia and provide it to the plant in return for reduced carbon generated via photosynthesis. This biological N₂-fixation provides a large proportion of the nitrogen in the natural environment (Fowler et al., 2013) and is an important component of sustainable agricultural systems, reducing the requirement for expensive nitrogen fertilizers and the pollution that can arise from their overuse (Vance, 2001).

The establishment of this symbiosis involves signaling between the two partners and results in rhizobia moving through an infection thread derived from an invaginated root cell wall into the root cortex where a new organ, the nodule is initiated. The cell wall of the root cells is degraded and the rhizobia released into the cell. Within the nodule infected cells, the rhizobia are enclosed in a plant-derived membrane to form an organelle-like compartment called the symbiosome. Within this symbiosome the rhizobia differentiate into their symbiotic form, the bacteroid. The symbiosome membrane (SM), initially derived from the plasma membrane (PM), becomes specialized as an interface between the bacteroid and its plant host, segregating the bacteroids from the plant cytoplasm and “protecting” them from any plant defense response (Mohd-Noor et al., 2015).

The major metabolite exchange across the SM is fixed nitrogen (principally ammonia) to the plant and a carbon source, most likely malate, to the bacteroids. However, transport of many other compounds into the symbiosome across the SM must occur as the enclosed bacteroids depend on the plant for all of their nutrients, including iron, zinc, calcium and cobalt amongst others (Brear et al. 2013; Udvardi and Poole, 2013; Clarke et al., 2014). The SM effectively controls the symbiosis via a suite of transport proteins synthesized by the plant. The plant can control what moves into the symbiosome and, presumably, can withhold sustenance if required. It has been suggested that the plant can impose sanctions on non-fixing rhizobia (Kiers et al.,

2003) and controlling transport across the symbiosome membrane could regulate this. It is also probable that compounds other than ammonia/ammonium move from the bacteroids to the plant (Udvardi and Poole 2013).

Transport studies with isolated symbiosomes have demonstrated the presence of a malate transporter and an ammonium channel on the SM, as well as metal ion transporters, but their molecular identity remains elusive (Udvardi and Day, 1997; Udvardi and Poole 2013; González-Guerrero et al. 2016). A number of proteomic analyses of the SM have been reported (Wienkoop and Saalbach, 2003, Catalano et al. 2004, Clarke et al. 2015), and although the earlier studies were limited by the lack of genome sequences for the legumes studied, an array of putative transport proteins have been identified. An example is LjSST1, a sulphate transporter later shown to be essential for nitrogen fixation in the *Lotus japonicus*- rhizobia interaction (Krussell et al. 2005; Schneider et al. 2019).

The most recent analysis of the soybean SM proteome (Clarke et al. 2015) identified a protein from the Yellow Stripe-like (YSL) family, Glyma.11G203400 (known then as Glyma11g31870). YSL proteins are members of the wider oligopeptide transporter family, generally considered to transport metals chelated to phytosiderophores (PS), such as deoxymugineic acid and nicotianamine (NA) (Curie et al., 2009). In monocots, PS excreted to the rhizosphere chelate ferric iron, and the complexes are transported into the plant cytoplasm by YSL transporters. Maize mutants for the first characterized member of this family show a phenotype of interveinal chlorosis that is characteristic of iron deficiency, and it is this phenotype that gave rise to the name Yellow-stripe 1 (YS1). YSL proteins, often localized in xylem parenchyma, can also transport other metal chelates (Dai et al. 2018; Chu et al., 2010; Sasaki et al., 2011, Zheng et al. 2012) and are important for intracellular iron transport and iron homeostasis, with both ferric and ferrous-PS complexes transported (Lubkowitz, 2011). YSL proteins are also involved in mobilization of intracellular stores of metals (Divol et al. 2013, Conte et al. 2013). YSL transporters operate through proton co-transport driven by the membrane potential (Schaaf et al. 2004), and whether localized to the PM or internal membranes, transport is always into the cell cytosol (Lubkowitz 2011).

Despite the biochemical characterization of some members of the YSL family, the functional role of other members is less clear. In particular, members of one phylogenetic clade of the YSL family, including Yellow Stripe-like 5, 7 and 8 (Group III) are not well characterized. A recent study showed that *Arabidopsis* YSL7 and 8 are responsible for the import of a *Pseudomonas syringae* virulence factor, syringolin A (Syl A), into the plant cytoplasm (Hofstetter et al. 2013). Syl A is a peptide derivative that acts by inhibiting the plant proteasome, and peptides of 4-8 amino acids in length inhibited its transport in plants and in yeast expressing AtYSL7. Consequently, it was suggested that AtYSL7 and AtYSL8 act as oligopeptide transporters, although direct evidence of oligopeptide transport was not shown (Hofstetter et al. 2013).

In this study we show that both AtYSL7 and soybean Yellow Stripe-like 7 (GmYSL7, encoded by Glyma.11G203400) can transport oligopeptides and that the soybean protein, which is localized to the SM in nodule infected cells, is important for nitrogen fixation.

RESULTS

GmYSL7 is a transporter of the YSL family

In our proteomic study (Clarke et al., 2015) we identified Glyma.11G203400 on the SM of soybean nodules. The protein is a member of the oligopeptide transporter (OPT) superfamily (Saier, 2000; Yen et al., 2001; Stacey et al., 2008) and has significant homology with members of the YSL family (Curie et al. 2009). We named it YSL7, as its closest *Arabidopsis* homologue is AtYSL7 (74% amino acid identity and 85% similarity; Yorden et al. 2011).

GmYSL7 belongs to a family consisting of 15 members in soybean (Supplemental Fig. S1; Schmutz et al., 2010) which in phylogenetic analysis fall into the three clades with both monocots and dicots members (Groups I – III, Supplemental Fig. S1). Group IV has only monocot members. In Genbank six proteins are annotated as “probable metal-nicotianamine transporter YSL7” but in the phylogenetic analysis only GmYSL7, Glyma.11G203400, associates closely with AtYSL7 in Group III, also clustering with the chickpea protein CaYSL7 (Ca08876) and three *M. truncatula* proteins (Medtr3g063490 [MtYSL7], Medtr3g063520 [MtYSL9] and Medtr5g091600

[MtYSL8]) (Supplemental Fig. S1). Of the other soybean proteins annotated as YSL7, Glyma.09G164500 and Glyma.16G212900 are more closely related to AtYSL5 and AtYSL8, while Glyma.09G281500, Glyma.20G004200 and Glyma.20G004300, although part of Group III, form a sub-clade not associated with any YSL proteins from other plants included in the phylogeny (Supplemental Fig. S1).

GmYSL7 is expressed in infected cells of soybean root nodules

Publicly available transcriptomic data for soybean suggests nodule-specific expression of *GmYSL7* (Severin et al. 2010; Supplemental Fig. S2). We confirmed this by measuring *GmYSL7* transcript abundance in leaves, roots of 8-day old seedlings, nodules and denodulated roots of 32-day-old plants using reverse transcription quantitative PCR (RT-qPCR). *GmYSL7* transcript was abundant in nodules but almost undetectable in other plant organs examined (Fig. 1A). We investigated the expression patterns of other YSL genes and all had lower nodule expression than *YSL7* and transcripts present in other tissues (Supplemental Fig. 2B).

Expression of *GmYSL7* during nodule development was examined. Transcripts were almost undetectable in young (6 – 10-day-old) inoculated roots, but abundance increased sharply before nitrogenase activity was first detected (day 18). Expression peaked in nodules of 18-day-old plants, and steadily decreased in older nodules (Fig. 1B; Supplemental Fig. 3).

As some characterized YSL proteins are involved in transport of iron complexes we examined expression of *GmYSL7* in nodules grown in hydroponics with a range of iron concentrations (0 – 100 μ M). *GmYSL7* expression was largely insensitive to iron concentration (Supplemental Fig. 3A). This was in contrast to two *AtYSL3* homologues with clear upregulation in high iron conditions (Supplemental Fig. 3 B, C).

We investigated *GmYSL7* cellular expression pattern in nitrogen-fixing nodules. The 2 kb genomic fragment immediately upstream of the coding region of *GmYSL7* was inserted upstream of a promoter-less green fluorescent protein- β -glucuronidase (GFP-GUS) coding sequence to give *pGmYSL7:GFP-GUS*. GUS staining of *pGmYSL7:GFP-GUS* transformed roots and nodules agreed well with our RT-qPCR

data, with no staining detectable in roots or in early nodule initials (Fig. 1C). GUS staining became evident as nodules developed (Fig. 1D) and was strongest in maturing nodules. In mature nodules, GUS staining was detected in the infected region and appeared to be confined to rhizobia-infected cells (Fig. 1E and F). No GUS staining was detected in the outer cortex of the nodule (Fig. 1E).

Localization of GmYSL7 on the SM in rhizobia-infected cells was confirmed using transgenic nodules expressing *pGmLbc3:GFP-GmYSL7* and analysed by confocal microscopy. FM4-64, a lipophilic dye that fluoresces when bound to membrane (Vida and Emr, 1995), was used to counterstain the SM (Limpens et al. 2009, Gavrin et al. 2014). GFP-GmYSL7 signal was on internal membranes within infected cells but not on the PM (Fig. 2A, B). Co-localization of GFP-GmYSL7 and FM4-64 signals was analysed by JACoP and Coloc 2 ImageJ plugins and quantified as Overlap coefficient with standard error (0.87 ± 0.05 ; Fig. 2C, D). Discrete localization of GFP-GmYSL7 on the SM can be seen also in Fig 2E, with GFP-YSL forming a clear “halo” around the perimeter of symbiosomes. The GFP-YSL7 fluorescence pattern in infected cells was distinct from free GFP, detected in the cytoplasm (Fig. 2F), and from that of a construct targeted to the symbiosome space (Fig. 2G).

Further confirmation that GmYSL7 was localized on the SM and not the PM was obtained by proteomic analysis of isolated SM and microsomal extract enriched in PM and endoplasmic reticulum. Approximately six times more peptides from the well characterized, SM-localized GmNOD26 were in the SM sample compared to the microsomal membrane sample, indicating enrichment of the SM in the purified sample. GmYSL7 peptides were only in the purified SM sample (Supplemental Table 1).

Silencing of GmYSL7 interrupts development of the symbiosis

Since GmYSL7 is localized to the SM, we investigated whether it is important for development of the symbiosis and nitrogen fixation by rhizobia using RNA interference (RNAi). Nodules from transgenic roots silenced for GmYSL7 were analyzed 24 days post inoculation (24 dpi; Fig. 3). Expression of *GmYSL7* in RNAi nodules was approximately 40% of the control but expression of the closest homologs, *Glyma.16G212900* (*GmYSL8*) and *Glyma.09G164500* (*GmYSL5*), was not

affected (Fig. 3A). Acetylene-reduction analyses showed that nitrogenase activity was reduced in silenced nodules to only 25% of the activity of control nodules (Fig. 3B). *GmYSL7* silenced nodules were smaller (Fig. 3C), paler (Supplemental Fig. S4 A-C) and displayed a delay in development (Fig. 3G-I) in comparison to empty vector control nodules (Fig. 3 D-F).

Silencing of *GmYSL7* did not affect bacteria release, but infected cells remained small and, unlike the control nodules, contained small, single-bacteroid symbiosomes (Fig. 3I). Numerous small vacuoles were localized around the nucleus (Fig. 3H) whereas control nodules had no vacuoles (Fig. 3E). To pinpoint the developmental stage in wild type nodules that matches the RNAi nodules we completed an analysis of nodules from soybean infected with *Bradyrhizobium diazoefficiens* strain 1042-45 carrying the *lacZ* fusion driven by the *nifD* promoter (Acuña et al., 1987). Four stages of development were identified and images can be seen in Supplemental Fig. S5. The morphology of infected cells of *GmYSL7*-silenced nodules (Fig. 3G-I) appeared to be arrested at stage II of normal nodule development (Fig. 3D-F, Supplemental Fig. 5E-H) where numerous small vacuoles were present in infected cells and most symbiosomes contained single elongated bacteroids. Electron microscopy (EM) of the silenced nodules confirmed that the infected cells were small and under-developed, packed with symbiosomes containing only a single bacteroid (Fig. 3I). Bacteroids appeared elongated as seen in control nodules during stage II (Supplemental Fig. 5F, G). Infected cells also contained numerous vacuoles of different sizes and apparent endosomes fusing with symbiosomes, reminiscent of the formation of a lytic compartment, which usually occurs during nodule senescence (Fig. 3I). Symbiosomes were isolated from silenced and control nodules, and this showed that in the silenced nodules, symbiosomes contained only single bacteroids compared to the control symbiosomes, which had multiple bacteroids (Supplemental Fig. 4D), confirming the phenotype seen by EM analysis. The results suggest that silencing of *GmYSL7* arrests development of soybean nodules at stage II.

RNAseq of nodules in which *GmYSL7* is silenced

We used RNAseq to compare the transcriptome in 22-day old nodules from *GmYSL7*-RNAi plants and empty vector controls. Before library construction, the level of *GmYSL7* transcript in each RNAseq sample was measured by RT-qPCR and silencing

of *GmYSL7* was confirmed in each of the *GmYSL7* RNAi samples (Supplemental Fig. 6). Principal component analysis (Fig. 4A) and a heatmap of gene expression of all differentially expressed genes shows clear differences between the RNAi and control samples (Fig. 4B). There were no significant changes in expression of other YSL genes.

There were 924 genes with \log_2 fold change of 1 or greater in nodules in which YSL7 was silenced, while 1180 genes had \log_2 fold change of -1 or greater (with adjusted p-value <0.05). Gene ontology (GO) enrichment analysis showed that genes involved in defence responses (e.g. defence response to bacterium, defence response to other organism), “negative regulation of endopeptidase activity” and a network associated with iron homeostasis, sequestration and transport, are overrepresented in the upregulated genes (Fig. 5A). A network of genes with GO terms including regulation of defence response, regulation of jasmonic acid mediated signaling pathway and regulation of signal transduction, and another including those associated with lipid biosynthesis, are overrepresented in the downregulated transcripts (Fig. 5B).

Details of expression of all genes in the RNAi and control nodules are available in Supplemental Table 2. Among the genes with significantly higher expression in the silenced nodules were those encoding homologues of a senescence-associated gene 13 (Glyma.12G059200), NRT1.5 (nitrate transporter 1.5, Glyma.18G260000), organic cation/carnitine transporter4 (Glyma.12G216400), ferritin (Glyma.01G124500, Glyma.11G232600), vacuolar iron transporter-like proteins (Glyma.05G121200, Glyma.08G076000), plantacyanin (Glyma.08G128100), a copper transport protein (Glyma.09G179800), cation efflux family protein (Glyma.08G164800), a cationic amino acid transporter 2 (Glyma.19G116500), nitrate transporter 2.4 (Glyma.11G195200) and a number of protease inhibitors (Supplemental Table 2).

Genes with significantly lower expression in the *GmYSL7* silenced nodules were those encoding homologues of sucrose-proton symporter 2 (Glyma.16G156900), glutamine dumper 2 (Glyma.18G277600), a number of carboxyesterases (alpha/beta-Hydrolases or carboxyesterase 17, Glyma.12G096400.1, Glyma.18G298000, Glyma.08G364100, Glyma.04G081600), Protein TIFY 5A-Related (Glyma.20G065500), nicotianamine synthase (MtNAS2 homologue, Glyma.15G251300), nodulin MtN21/EamA-like

transporter family protein (Glyma.06G246000), cytochrome P450, family 94, subfamily C, polypeptide 1 (Glyma.11G212000) and GmNIC1a (Glyma.12G208900) a Clavata3/ESR (CLE) related homologue (Supplemental Table 2).

GmYSL7 transports oligopeptides and Syringolin A but not Fe(II)-NA

Yeast complementation was used to try to identify a substrate for GmYSL7. Initially we tested for transport of Fe(II)-NA by complementation of the *fet3/fet4/ftr1* mutant; however, although the positive control *ZmYSI* (Curie et al. 2001, Schaaf et al. 2004) complemented the mutant, *GmYSL7* and *AtYSL7* did not (Fig. 6).

Since the YSL family is part of the wider oligopeptide transporter (OPT) family, we next tested whether GmYSL7 could complement the yeast oligopeptide transport *opt1* mutant, using different oligopeptides as the sole source of nitrogen for growth. The peptides chosen included peptides used in other yeast assays for transport of oligopeptides (Osawa et al. 2006), peptides used in Hofstetter et al. (2013) and a peptide associated with a biological role in nodules (CLE peptide, GmRIC1a encoded by Glyma.13G292300; Hastwell et al. 2015). When the transformants were grown with four (ALAL, LSKL), five (IIGLM) and six (KLLLLG) amino acid peptides as the only N source, cells expressing *AtYSL7*, *GmYSL7* or *AtOPT4*, but not with the empty vector pDR196, grew (Fig. 7). On media containing larger peptides (eight, ten or twelve amino acid), the growth of the transformants was more varied. *AtOPT4* supported growth on the eight amino acid peptide DRVYIHPF, while growth was weak for *AtYSL7* and *GmYSL7*. Growth on the 10 amino acid peptides DRVYIHPFHL was close to background for all transformants, but all grew better than vector control on the 12 amino acid peptide RLAPGPDPHH (GmRIC1a, Fig 7).

AtYSL7 is involved in syringolin A (SylA) uptake and, when expressed in yeast, exposure to SylA inhibited growth (Hofstetter et al. 2013, Fig. 8) suggesting the transporter mediated uptake of this toxic peptide derivative. We used this assay to test for transport of SylA by GmYSL7. *GmYSL7*, *AtYSL7* and the empty vector pDR195 were expressed in the yeast *pdr5* mutant that lacks the ABC transporter PDR5, and plated as a lawn (Hofstetter et al., 2013). When a disk containing SylA was placed on the plate, growth of yeast expressing *GmYSL7* and *AtYSL7*, but not the empty vector pDR195, was inhibited. Inhibition of growth caused by SylA on the *AtYSL7* plate

showed as clear patches on the plate for all concentrations of SylA tested, while inhibition of yeast expressing GmYSL7 was weaker (Fig. 8).

GmYSL7 can replace MtYSL7

MtYSL7 (*Medtr3g063490*) is the closest homolog of GmYSL7 in *Medicago truncatula*, although it is localized on the PM in the vasculature and nodule cortex (Castro-Rodríguez et al. 2020). To determine whether GmYSL7 and MtYSL7 proteins play similar roles in the different cell types in which they are located, we expressed *GmYSL7* in the *Mtysl7-1* mutant (Castro-Rodríguez et al. 2020). Expression was driven by the *MtYSL7* promoter to ensure expression in the cells in which MtYSL7 is present (vasculature and nodule cortex but not infected cells). Although GmYSL7 localizes to the SM in soybean, it was able to complement the *Mtysl7-1* transposon insertion mutant to restore nitrogenase activity to wild type levels and increase the dry weight of the transformed plants compared to the mutant (Fig. 9). Based on this result, we assume that in the *Mtysl7-1* mutant, when expressed in the cells where MtYSL7 is normally active, GmYSL7 at least partially localizes to the PM.

DISCUSSION

We have characterized a member of the YSL family, GmYSL7, in soybean. The protein is part of a clade of YSL proteins (Group III) that includes AtYSL5, AtYSL7 and AtYSL8. AtYSL7 and 8 are involved in transport of the *Pseudomonas syringae* virulence factor into Arabidopsis cells across the PM (Hofstetter et al. 2013), but their physiological role in plants has not been determined. GmYSL7, AtYSL7, CaYSL7 and three *Medicago truncatula* proteins (MtYSL7, 8 and 9) form a cluster in phylogenetic analyses, but the soybean protein's expression profile is distinct from that of *AtYSL7* and *MtYSL7*. *AtYSL7* is expressed mainly in flowers but also in siliques and roots. *MtYSL7*, has highest expression in nodules, but is also expressed in roots. Soybean, on the other hand, appears to lack an *YSL7* paralog with expression similar to *AtYSL7* and *MtYSL7*. Rather, *GmYSL7* expression is linked specifically to symbiotic nitrogen fixation, occurring only in infected nodule cells where the protein is present on the SM, but not the PM, in contrast to *AtYSL7* and *MtYSL7*. Furthermore, its expression is only marginally affected by the iron concentration of the growth medium (Supplemental Fig. 3). This seems a clear example of

neofunctionalization with the loss of the paralog (Xu et al. 2017). Either there is no requirement in soybean for the role played by AtYSL7 in other organs or another soybean gene with functional redundancy fulfills that role. The closest homologues of *GmYSL7*, *Glyma.16G054200* (*GmYSL8*) and *Glyma.19G094800* (*GmYSL5*) are expressed in almost all tissues (Supplemental Fig. 2), but we know nothing about their function at this stage.

An important role for *GmYSL7* in nitrogen-fixing nodules is shown by knockdown of its expression, which resulted in smaller nodules, with reduced leghemoglobin and a decrease in nitrogenase activity. The *GmYSL7*-RNAi nodules appear to have been developmentally arrested, with infected cells having small single bacteroid symbiosomes, in contrast to the large symbiosomes containing multiple bacteroids in infected cells of control nodules. The *GmYSL7*-RNAi infected cell ultrastructure is similar to control nodules in the early stages of development. This suggests that the activity of *GmYSL7*, and the substrate(s) it transports across the SM, is important for the continued development of the symbiosis and maturation of infected cells.

Our results show clearly that *GmYSL7* transports a range of small peptides. When considering the activity of SM transporters, it is important to bear in mind the orientation and energization of the SM (Udvardi and Day, 1997), as this influences the direction that any given substrate is transported. A P-type ATPase on the SM together with the rhizobial electron transport chain pump protons into the symbiosome space, creating an electrochemical gradient across the SM, with the membrane potential positive on the inside and acidifying the interior of the symbiosome (symbiosome space or peribacteroid space) (Udvardi and Day, 1997). All YSL proteins characterised to date transport compounds across cell membranes into the cytoplasm (Lubkowitz 2011), with proton symport the most likely mechanism (Schaaf et al. 2004). Assuming that *GmYSL7* has a similar mechanism, it is consequently likely to transport its peptide substrate out of the symbiosome and into the plant cell cytosol. The phenotype seen in *GmYSL7*-RNAi nodules is, therefore, related to the lack of provision of this substrate to the plant cell.

A *Medicago truncatula* homologue of *GmYSL7*, *MtYSL7*, has been characterized and also transports oligopeptides. It is not localized on the SM but the *Mtysl7*-mutant

has a phenotype that also affects the symbiosis and nitrogen fixation (Castro-Rodríguez et al. 2020). The difference in cellular localization might be explained by the fact that *M. truncatula* produces indeterminant nodules, where the meristem continues to be active throughout development, and soybean determinant nodules, in which mature nodules have no meristem. As well as structural differences the different nodule types have a number of metabolic differences including in the mechanism for nitrogen assimilation and the compounds transported from the nodules (Smith and Atkins, 2002). However, *GmYSL7* was able to complement the *Mtysl7-1* mutant, restoring nitrogenase activity and growth in low N conditions. This suggests that the two YSL7 proteins are able to transport the same substrate/s and that while MtYSL7 brings this substrate into the cell across the PM, GmYSL7 moves its substrate out of the symbiosome and into the cytosol.

In contrast to many other YSL proteins, GmYSL7, AtYSL7 and MtYSL7 were not able to transport Fe(II)-NA (this study, Castro-Rodríguez et al. 2020). Additionally MtYSL7 could not transport Fe(III), zinc or copper complexed with NA (Castro-Rodríguez et al. 2020). On the other hand, complementation of the yeast *opt1* mutant showed that the three proteins could transport oligopeptides of various sizes and amino acid sequences. Like AtYSL7, when expressed in yeast, GmYSL7 could also transport syringolin A, a peptide derivative that is the virulence factor for *Pseudomonas syringae* (Hofstetter et al. 2013). Inhibition of yeast growth caused by the transported Syla was not as strong as for AtYSL7 suggesting that GmYSL7 may not transport it as effectively or have the same specificity for the compound. In our assays for direct uptake of oligopeptides in yeast, both AtYSL7 and GmYSL7 supported growth on media with oligopeptides of 4-6 and 12 amino acids as their sole N source, but there was little growth when the 8 amino acid peptide, DRVYIHPF was used, despite its ability to reduce the effect of Syla on Arabidopsis roots (Hofstetter et al., 2013).

While it is clear that YSL7 proteins are peptide transporters, their physiological role in legumes is not clear. Glutathione is a three amino acid peptide derivative found in nodules and bacteroids but as MtYSL7 cannot transport GSH (Castro-Rodríguez et al., 2020) and GmYSL7 is able to replace the function of MtYSL7, it is unlikely that transport of GSH out of the symbiosome is the physiological role of GmYSL7. The

symbiosome contains a number of proteases on the SM and in the symbiosome space (Clarke et al. 2015) and appears to act like a vacuole containing large amounts of free peptides (Clarke et al. 2015). Some of these could be substrates for GmYSL7, but why blocking their exit from the symbiosome would inhibit N-fixation and symbiosome development to such an extent is not obvious. While it is possible that GmYSL7 acts to scavenge N by transporting peptides from the symbiosome space into the plant cytosol, it is unlikely that this would have such a profound effect on nodule development.

It is tempting to speculate that release of peptides from the symbiosomes has a more direct role in manipulating plant gene expression and organogenesis. Cyclic peptides act as signaling molecules in some symbioses (Abbamondi et al. 2014) and it is possible that GmYSL7 transports an oligopeptide derivative produced in the bacteroids. In this scenario, release of the oligopeptide signal could be required to relieve plant inhibition of bacteroid division or as a positive signal for symbiosome development. Supporting this idea is the fact that a protein annotated as an oligopeptide transporter was specifically induced in symbiotic *Bradyrhizobium japonicum* (Pessi et al. 2007) and a number of transcription factors are up- or downregulated in nodules of *GmYSL7*-RNAi plants.

We used RNAseq of *GmYSL7*-RNAi nodules to investigate further the effects of inhibiting transport by GmYSL7. Overrepresented GO terms among the downregulated genes included regulation of defense response and regulation of jasmonic acid signalling pathway and also a range of terms associated with lipid metabolic processes (lipid biosynthetic process, isoprenoid metabolic process). The lipid associated terms likely relate to the failure of the symbiosome to develop with multiple bacteroids. The change from a single bacteroid symbiosome to one with multiple bacteroids is likely to require synthesis of large amounts of lipid. With development of the infected cell blocked at an early stage this synthesis would not be required.

Overrepresented terms in the upregulated genes included “defense response”, “defense response to bacteria”, “defense response to other organism” and “regulation of defense response”, suggesting that blocking transport by GmYSL7 causes a general

defense response against the rhizobia. This links with the downregulation of genes associated with the regulation of defense response GO term which may indicate that the inhibition of the defense response that allows the symbiotic association to develop, has been lifted in the RNAi nodules. The changes in genes associated with a defense response may be an indirect effect of a decrease in nitrogen fixation, with the plant sanctioning the bacteria as ineffective (Kiers et al. 2003). Upregulation of Glyma.11G195200, a soybean homologue of AtNRT2.4, a nitrate transporter that is upregulated in response to nitrogen starvation, suggests that as nitrogenase activity was reduced, the nodules in *GmYSL7*-RNAi plants may indeed be nitrogen starved. A laser ablation electrospray ionization-mass spectrometry (LAESI-MS) study comparing ineffective *nifH* nodules with wild type (Agtuca et al., 2020) found jasmonic acid (JA) associated metabolites were increased in the nitrogen deprived *nifH* nodules. JA is known to mediate defense responses with an associated increase in secondary metabolites (Pauwels et al. 2009). Two of the transcripts with strongest downregulation, Glyma.20G065500.1 and .2, and a third, Glyma.11G212000, are likely to encode proteins that repress JA defense responses, which would likely lead to an increase in JA and initiation of a defense response to rhizobia.

Zeatin is another metabolite associated with functional nodules which has reduced abundance in non-fixing *nifH* nodules (Agtuca et al., 2020). In the *GmYSL7*-RNAi nodules, that also have reduced nitrogen fixation, a gene encoding isopentyl transferase 5 (*GmIPT5*, Mens et al. 2018) is downregulated. *GmNIC1*, a peptide responsible for nitrate regulation of nodulation, is also downregulated in the RNAi nodules indicating that there is no requirement for control of nodule numbers in the RNAi plants.

A link to peptide signalling in the RNAi nodules is the downregulation of three genes encoding glutamine dumpers (GDU). Glyma.18G277600 is one of the most downregulated genes in RNAi nodules with expression reduced 29-fold, while Glyma.14G105200 and Glyma.17G083000 are also reduced significantly. In *Arabidopsis*, most GDUs are regulated via a peptide receptor PEPR that interacts with *AtPep1* (Ma et al 2014). There is no orthologue of *AtPep1* in soybean but homologues of the PEPR gene (Glyma.20G194400 and Glyma.10G195700) are highly expressed in soybean nodules (this study) and could be involved in perception

of the peptide transported by YSL7. In Arabidopsis, glutamine dumpers are involved in regulating amino acid exporters via an interaction with a ubiquitin ligase (LOG2) (Guerra et al. 2017), homologues of which are expressed in nodules.

Although soybean exports most nitrogen from nodules as ureides, it also exports some amides (Smith and Atkins, 2002), so it is possible that the GDU downregulated in *GmYSL7*-RNAi nodules is involved in export of fixed nitrogen. A purine permease, likely required for uptake of these purines in the uninfected cells, is also downregulated in the RNAi nodules. It is possible, therefore, that the peptide transported via *GmYSL7* from the symbiosome provides a link between nitrogen fixation in the bacteroid and fixed-N transport from the nodule. Export of a peptide from the bacteroid to the plant that increases fixed-N transport out of the nodule, would prevent feedback inhibition of nitrogen fixation by the plant. Disruption of nitrogen fixation in the bacteroid, on the other hand, may inhibit peptide transport, decreasing fixed-N efflux from the nodule to maintain C/N balance in the nodule.

Another possibility is that GDU regulates export of amino acids across the symbiosome membrane. A number of studies suggest that the majority of the fixed nitrogen is exported and not assimilated by the bacteroids (reviewed by Patriarca et al. 2002); bacteroids down regulate glutamine synthetase and symbiotic auxotrophy for branched chain amino acids has been demonstrated in pea (Prell et al. 2009) and *Phaseolus vulgaris* (Prell 2010). In bacteroids of soybean nodules, genes encoding proteins associated with amino acid biosynthesis are downregulated when compared to free-living bacteria (Pessi et al. 2007), suggesting that plants may supply amino acids to support nitrogen fixation. As import into the symbiosome is equivalent to export from the cell this would require the action of an amino acid exporter. In this scenario export of the peptide that is the substrate for YSL7 would regulate GDU expression and the supply of amino acids to the symbiosome and bacteroid allowing continuation of the symbiotic interaction.

Another group of GO terms that are overrepresented in the upregulated genes in *GmYSL7*-RNAi nodules are “intracellular sequestering of iron ion”, “iron ion transport” and “cellular iron ion homeostasis”. Some of the genes associated with

these terms include ferritins and vacuolar iron transporter (VIT) gene homologues. This is accompanied by upregulation of a number proteins potentially associated with metal transport (Cu transport protein, cation efflux family protein, MATE efflux family protein) and a transcription factor, WRKY9, associated with a GO term “cellular response to iron ion starvation”, and downregulation of a gene encoding NAS, responsible for synthesis of NA, a PS involved in metal transport by YSLs. MtNAS2 is essential for iron supply to the nodules in *Medicago truncatula* and in the *nas2-1* mutant nitrogenase activity is reduced (Escudero et al. 2020). These results indicate that metal homeostasis is dysregulated in the *GmYSL7*-RNAi nodules. There are two possible explanations for this. *GmYSL7* may play a direct role in metal ion homeostasis with the peptide transported acting as a signal to regulate metal homeostasis. However, since the genes with modified expression are not those most highly expressed in nodules, we favor the second explanation, namely that when symbiosome development is stalled and nitrogen fixation blocked, then metals being supplied to the symbiosome by the plant may accumulate in the nodules and need to be sequestered to avoid cellular damage. This may result in storage of metals, particularly iron, in uninfected cells with ferritin, or transport into the vacuole via VIT proteins.

Further study is required to decide between these options. Identification of the peptide substrate for *GmYSL7* will be key to determining its role.

Conclusion

We have identified a member of the oligopeptide transporter family, *GmYSL7*, which is localized to the symbiosome membrane in nitrogen-fixing soybean nodules. It transports an array of small oligopeptides out of the symbiosome and into the plant cell cytosol, and its disruption arrests infected cell development and symbiosome maturation, inhibiting nitrogen fixation. *GmYSL7*'s disruption affects expression of a number of genes involved in plant defense responses, nitrogen metabolism and in metal homeostasis suggesting a role monitoring the functional state of the bacteroids and regulating nodule metabolism and transport processes. The symbiosome membrane localization has been ideal to illustrate the effects of oligopeptide transport by *GmYSL7* but the ability to transport oligopeptides is shared by *AtYSL7* and presumably other YSL7 proteins. It will be interesting in future to determine whether

the same substrates are physiologically important in different tissues and other plants and the roles this oligopeptide transport plays in plant growth and development.

METHODS

Plant Growth Conditions

Soybean (*Glycine max* L.) cv Stevens seeds were inoculated at planting and one week after planting with *Bradyrhizobium diazoefficiens* (Soybean group H, New Edge Microbials). Plants were grown as described in Clarke et al. (2015) and fertilized once a week with a nitrogen-free B&D nutrient solution (Broughton and Dilworth 1971). Nitrogenase activity in nodules was assessed using an acetylene reduction assay as described by Unkovich et al. (2008).

For limited and excess iron conditions plants were grown in B&D solution with 0, 1, 10 (control concentration) or 100 μ M Fe-citrate, which was renewed every 2 days to maintain pH and stable nutrient supply. Two biological replicates were done. Fe status was determined by elemental analysis (Lee M, School of Land and Environment, University of Melbourne) using the Perchloric Nitric Acid Method. 15 plants per treatment were analyzed to determine shoot iron content using an Inductively Coupled Plasma Optical Emission Spectrometer (Varian Medical Systems, Palo Alto, CA, USA).

Cloning and Constructs

Genomic DNA for cloning the *GmYSL7* promoter was extracted from mature soybean leaves using DNeasy Plant minikit (Qiagen). RNA was extracted from plant tissues using an RNeasy Plant mini kit (Qiagen) and cDNA synthesized using an iScript cDNA synthesis kit (Invitrogen). All constructs were PCR amplified from *Arabidopsis thaliana* seedling or soybean nodule cDNA, gDNA or available plasmids using either Platinum Pfx50 (Invitrogen) or Phusion (Thermo Fisher Scientific) high fidelity polymerases and cloned using the Gateway cloning system (Invitrogen) to produce entry clones (Brear et al. 2020) for GmYSL7, AtYSL7, ZmYS1 and AtIRT1. A list of primers used can be found in Supplemental Table 3. The entry clones were used as a basis for LR recombination to produce most expression constructs.

For GmYSL7 promoter GUS fusion constructs, a 2 kb genomic fragment immediately upstream of the GmYSL7 coding region was recombined into either pKGW-GGRR (Gavrin et al. 2016) or pKGWFS7 (Karimi et al., 2002). The full-length coding sequence of GmYSL7 was recombined into pGmLBC3-pK7GWIWG2 Gateway vector (Gavrin et al. 2016) to create a hairpin RNAi vector for silencing the gene. N-terminal GFP fusion constructs for GmYSL7 were constructed from the full-length coding sequence recombined into either pGmLBC3-pK7WGF2-R or a modified pK7WGF2 (pGmLBC3-pK7WGF2) where the 35S promoter is replaced by the GmLBC3 promoter (Gavrin et al. 2016). The free GFP construct was made by *EcoRV* digestion and re-ligation of the pGmLBC3-pK7WGF2 vector to remove the intervening Gateway cassette. The symbiosome space MtNOD25-GFP construct is from Hohnjec et al. (2009) and encodes the first 24 amino acids of MtNOD25 fused to GFP. For yeast expression, full length open reading frames of GmYSL7, AtYSL7, and AtIRT1 were recombined into the pDR196GW vector (Brear et al. 2020). AtOPT4 (Osawa et al. 2006) and ZmYS1 (Curie et al. 2001) in pDR196 (Rentsch et al. 1995) were provided by Doris Rentsch and Catherine Curie, respectively.

The GmYSL7 coding sequence was synthesised with the MtYSL7 promoter and flanked by attL recombination sites inserted in the pUC57 (Synbio). The construct was recombined into pGBW13 using Gateway Cloning technology.

Transformation of Soybean and Medicago

Hairy root transformation of soybeans (cv Stevens) used *Agrobacterium rhizogenes* K599 and was as described by Mohammadi-Dehcheshmeh et al. (2014). Transformed roots were inoculated with *B. diazoefficiens* CB1809 (Becker Underwood, Somersby, NSW, Australia). Plants were grown under controlled temperature and lighting conditions (26°C day, 24°C night; 16 hr day; 120-150 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Transformed nodules were examined 2-4 weeks post inoculation.

Transformation of *Medicago truncatula* was as described by Boisson-Dernier et al. (2001) using *A. rhizogenes* ARqua1.

Microscopy

Confocal imaging of GFP-fusion proteins was done on transgenic nodules either hand sectioned or sectioned in low melt agarose using a vibratome (752M Vibroslice, Campden Instruments, Loughborough, Leics., UK). In some instances, nodules were counterstained by FM4-64 (30 µg/ml). Nodule sections were immediately imaged as described previously (Limpens et al., 2009) using either an LSM Pascal 410 (Zeiss) or an SP5 II (Leica) confocal laser-scanning microscope equipped by an argon laser with a 488-nm laser line and helium-neon laser with a 543-nm laser line. Confocal settings: excitation at 488 nm for GFP and 543 nm FM4-64; GFP emission was selectively detected using a 505-530 nm range; FM4-64 emission was detected in another channel using a 560-615 nm range.

Imaging of GUS expression was done as described in Clarke et al. (2015). Sections were either counterstained with ruthenium red or mounted directly in Milli-Q water, and imaged using an Axiophot epifluorescence microscope with a set of Achromatic objective lens (Zeiss).

The protocol for tissue preparation for light and EM has been described previously (Limpens et al., 2009). Semithin sections (0.6 µm) for light microscopy and thin sections (60 nm) for EM of transgenic nodules were cut using a Leica Ultracut ultramicrotome UC7 (Leica). Sections were collected on 400 mesh nickel grids and examined using a Jeol JEM 1400 transmission electron microscope (Jeol Ltd, Tokyo, Japan).

Reverse Transcription Quantitative-PCR

RT-qPCR assays were used to measure transcript abundance in soybean tissues of control and YSL7 RNAi plants grown in sand or hydroponics. cDNA was synthesized from 500 ng total RNA using Iscript reverse transcriptase (Bio-Rad, Hercules, CA, USA), according to manufacturer's instructions. RT-qPCR assays were done in a volume of 5 µl in triplicate and contained 1 µl of cDNA diluted 1/5, 1 X LightCycler® 480 SYBR green I mix (Roche Applied Science, Castle Hill, Australia) and 0.5 µM of each primer (GmYSL7 and GmUBI3 QRT primers; Supplemental Table 3). Assays were done using a LightCycler® 480 (Roche Applied Science) and the following conditions: 95°C 10 min, 45 cycles of 95°C 10 s, 56°C 10 s, 72°C 20 s,

followed by ramping the temperature from 55°C to 95°C for melt curve analysis. PCR efficiency for each primer pair was determined using the LinRegPCR software (Ramakers et al., 2003) and data analysed using the LightCycler® 480 software package (Roche Applied Science). Data were normalized using *GmUBI3* (Glyma20g27950; Trevaskis et al., 2002) or *cons6* expression (Libault et al., 2008). Stable *GmUBI3* expression in the tissues examined in this study was confirmed through comparison of its expression with five characterized soybean reference genes (*cons4*, 6, 7, and 15; Libault et al., 2008) using geNorm software (Vandesompele et al., 2002). The amplified product from the real-time reaction was cloned and sequenced to confirm the specificity of the amplification product.

Yeast complementation

To test for transport of Fe(II)NA AtYSL7, GmYSL7, ZmYS1 in pDR196GW and the empty vector were introduced into the *Saccharomyces cerevisiae* (yeast) *fet3/fet4/ftr1* mutant (Spizzo et al. 1997; DEY1530: *MATa ade2 his3 leu2 lys2 trp1 ura3 fet3-2::HIS3 fet4-1::LEU2 ftr1D1::TRP1*) using the method described by Dohmen et al. (1991). Fe(II)-NA plates were prepared by mixing 15 µl 10 mM FeSO₄ in 200 mM MES/Tris pH 7.4 with 250 µl 200 mM Na-ascorbate and 8 µl of 50 mM NA and heating at 65°C for 10 minutes to produce a clear solution that was added to 25 ml of SD media to produce the Fe(II)-NA plate. Transformants were grown in liquid media to an OD₆₀₀ of 1 and then serially spotted in ten-fold dilutions on either SD plates with no added iron, Fe(II)-NA plates or SD-plates with 10 µM Fe-citrate.

For the peptide transport assay AtOPT4, AtYSL7, GmYSL7 in pDR196GW and the empty vector were introduced into the yeast *opt1* mutant (Y11213: BY4742; *MATa; ura3Δ0; leu2Δ0; his3Δ1; lys2Δ0; YJL212c::kanMX4*, Euroscarf). Transformants were grown as above on minimal medium (0.17% YNB without amino acids and (NH₄)₂SO₄, supplemented with amino acids as required, and containing either 10 mM NH₄Cl (positive control) or 100 µM of the following peptides, ALAL, LSKL, IIGLM, KLLLLG, DRVYIHPF, DRVYIHPFHL or RLAPGPDPPHHN, as the sole source of nitrogen.

Syringolin A transport assay

An assay for transport of Syringolin A by AtYSL7 and GmYSL7 in the yeast strain $\Delta pdr5$ (Y12409: BY4742; *MATa*; *ura3Δ0*; *leu2Δ0*; *his3Δ1*; *lys2Δ0*; *YOR153w::kanMX4*, Euroscarf) was done as described in Hofstetter et al. (2013). Syringolin A was kindly provided by Robert Dudler, University of Zurich.

Statistical Analyses

A one-way ANOVA with Tukey's HSD (SAS Enterprise Guide Version 4.3; SAS Institute Inc., Cary, NC, USA) was used to analyze differences in plant organ dry mass after growth in varying Fe concentrations. Differences are reported as significant where $p < 0.05$.

RNAseq analysis of transcriptome in *GmYSL7*-RNAi nodules

The transcriptome for *GmYSL7*-RNAi nodules was compared to those transformed with an empty vector control using RNAseq. Hairy root transformation with pGmLBC3-pK7GWIWG2 vector or the vector containing *GmYSL7* coding sequence was used to produce transformed nodules. RNA was isolated from nodules 21 or 22 days after inoculation using an RNeasy kit (Qiagen). Five replicates for each construct were done, each with nodules from 5-7 transformed plants. RNA integrity number (RIN) was determined on a 2100 bioanalyzer (Agilent) and was between 7 and 8.3 for all samples. RNAseq library construction and analysis were completed at Institute for Molecular Bioscience Sequencing Facility, The University of Queensland. A combination of the Ribo Zero rRNA removal bacteria (Illumina) and Ribo Zero rRNA removal plant (seed and root) (Illumina) was used to eliminate the rRNA from the sample. The library was constructed using a TruSeq® Stranded mRNA LT - SetA and SetB (Illumina). Sequencing was performed using the Illumina NextSeq500 (NextSeq control software v1.4/ Real Time Analysis v2.1). The library pool was diluted and denatured according to the standard NextSeq protocol, and sequenced to generate single-end 76 bp reads using a 75 cycle NextSeq500/550 High Output reagent Kit (Illumina).

Raw sequence reads were aligned to the JGI Wm82.a2 soybean assembly. DESeq2 (Love et al. 2014) was used to test for differential expression between control and YSL7-RNAi samples. Genes with \log_2 fold change (\log_2 FC) >1 and adjusted p-value < 0.05 were considered differentially expressed. Overrepresented biological terms

were identified from the list of differentially expressed genes. GO term enrichment analysis was based on the information in SoyBase (<https://soybase.org/>). Enriched biological terms and their linkage were analysed and visualized using ClueGO v2.5.5 (Bindea et al., 2009), implemented in the Cytoscape v3.5.1 environment (Shannon et al., 2003; <https://cytoscape.org/cy3.html>). ClueGO parameters were as follow: Analysis Mode, Functional Analysis; Load Markers List, *Glycine max* (3847); Visual Style: Significance Shape, ellipse; ClueGO settings, Ontology/Pathway; GO, Biological Process / KEGG: downloaded the 12/12/2017; Evidence type, All Evidences; Statistical Test Used = Enrichment/Depletion (Two-sided hypergeometric test); Correction Method Used = Bonferroni step down; Min GO Level = 3, Max GO Level = 8, Min Percentage = 4.0, GO Fusion = true, GO Group = true, Kappa Score Threshold = 0.4; Over View Term = SmallestPValue; Group By Kappa Statistics = true; Initial Group Size = 1; Sharing Group Percentage = 50.0.

Symbiosome isolation, SM and microsomal membrane isolation and proteomic analysis.

Symbiosomes were isolated as described by Clarke et al. (2015). SM was collected after pelleting of the membrane and resuspended in 1M Urea for proteomic analysis. Microsomal membrane was isolated from nodules ground and filtered through miracloth as described in Clarke et al. (2015). Symbiosomes and other intact organelles were pelleted by centrifugation at 20000 g. The membrane in the supernatant (enriched in PM and endoplasmic reticulum) was collected by centrifugation at 100,000 g for 1 hour at 4°C and the pellet resuspended in 8M Urea. Proteomic analysis was completed at the La Trobe Comprehensive Proteomics Platform (La Trobe University). Data were collected on a Q Exactive HF (Thermo-Fisher Scientific) in Data Dependent Acquisition mode using m/z 350–1500 as MS scan range at 60 000 resolution. HCD MS/MS spectra were collected for the 7 most intense ions per MS scan at 60 000 resolution with a normalized collision energy of 28% and an isolation window of 1.4 m/z . Dynamic exclusion parameters were set as follows: exclude isotope on, duration 30 s and peptide match preferred. Other instrument parameters for the Orbitrap were MS maximum injection time 30 ms with AGC target 3×10^6 , MSMS for a maximum injection time of 110 ms with AGT target of 1×10^5 .

Raw files consisting of high-resolution MS/MS spectra were processed with MaxQuant version 1.5.5.1 to detect features and identify proteins using the search engine Andromeda. Sequence data for soybean from Phytozome (https://phytozome.jgi.doe.gov/pz/portal.html#!info?alias=Org_Gmax) was used as the database for the search engine.

ACCESSION NUMBERS

The accession number for *GmYSL7* is NM_001289202.2

SUPPLEMENTAL DATA

Supplemental Table S1. Unique *GmYSL7* and *GmNOD26* peptides¹ identified in purified symbiosome membrane, a microsomal membrane fraction, and a symbiosome-enriched membrane sample from soybean nodule homogenate.

Supplemental Table S2. Genes upregulated or downregulated in *GmYSL7*-RNAi nodules and data for all genes expressed in the nodules.

Supplemental Table S3. Primers used in this study.

Supplemental Figure S1. Phylogenetic analysis of YSL proteins

Supplemental Figure S2. Expression analysis of the *YSL* genes in soybean.

Supplemental Figure S3. Expression of *GmYSL7* and two *YSL3* homologues in response to Fe status during nodule development.

Supplemental Figure S4. Physical characteristics of nodules and FM 4-64 stained symbiosomes extracted from empty vector control and *GmYSL7*-RNAi nodules.

Supplemental Figure S5. Detailed morphological analysis of determinate nodule development

Supplemental Figure S6. Expression level of *GmYSL7* in RNAseq samples measured by RT-qPCR.

ACKNOWLEDGMENTS

We thank Catherine Curie for providing the plasmid containing ZmYS1 and useful discussions about YSL transporters, Sarah Conte and Elsbeth Walker for providing advice about the methods for yeast assays for transport of Fe(II)-NA and Robert Dudler for providing Syringolin A.

FIGURE LEGENDS

Figure 1. GmYSL7 is expressed in infected cells of soybean root nodules. A. Transcript level of *GmYSL7* in tissue samples from different organs. DNR, denodulated roots; nd, not detected. B. Transcript level of *GmYSL7* during nodule development. 6R, roots 6 days after inoculation (DAI); 10RN, roots and nodules 10 DAI; 13N-26N, nodules the indicated DAI. Data shown are from three independent time courses. Error bars are SE (n = 3). Nitrogenase activity was first detected at 18 DAI. The relative expression was quantified by RT-qPCR and normalized to a soybean ubiquitin gene (*GmUBI3*, Glyma20g27950.1). C. Transgenic root expressing pYSL7:GFP-GUS. GUS staining was not detectable in the very early stages of nodule development. Arrowhead indicates a nodule initiation. D. Transgenic pYSL7:GFP-GUS 10-day-old nodule primordia. IR, infected region. E. Transgenic pYSL7:GFP-GUS mature nodule. GUS staining is restricted to infected cells. OC, outer cortex. F. Magnification of E. Scale bars, 150 μ m.

Figure 2. Localization of GmYSL7 in soybean nodule cells infected with rhizobia. A. Localization of GFP-GmYSL7 on symbiosome membranes in an infected cell counterstained by membrane lipophilic dye FM4-64; IC, infected cell; NI, non-infected cell; ROI, region of interest. B. Fluorescent intensity plot of ROI from A. C. Colocalization analysis of GFP-YSL7 with FM4-64. The colocalised points are highlighted by white in colocalization map. D. Scatterplot between the intensities of the GFP-YSL7 and FM4-64 in C. E. Superimposed confocal z-stack sections of GFP-GmYSL7 signal in rhizobia-infected nodule cells. F. Free GFP localizes to the cytoplasmic spaces surrounding symbiosomes in infected cells. G. MtNOD25-GFP (Hohnjec et al. 2009) localizes to the peribacteroid space inside the symbiosomes. Scale bars, 5 μ m.

Figure 3. RNAi silencing of *GmYSL7* affects nodule development. A. Transcript level of *GmYSL7* and its closest homologs in 24-day-old nodules of empty vector control and *GmYSL7*-RNAi plants (error bars represent SD; n=4; t-test: *, p<0.05). B. Nitrogenase enzyme activity in 24-day-old nodules of empty vector control and *GmYSL7*-RNAi plants (error bars represent SD; n=8; t-test: **, p<0.01). C. Fresh weight of 24-day-old nodules of empty vector control and RNAi-*GmYSL7* (error bars represent SD; t-test: *, p<0.05). D. Longitudinal section of a 24-day-old nodule from an empty vector control plant. Scale bar, 200 μ m. E. Magnification of (D) showing developed (stage IV) infected cells. Scale bar, 10 μ m. F. Electron microscopy of infected cells of a 24-day-old nodule from empty vector control containing developed multibacteroid symbiosomes. Scale bar, 2 μ m. G. Longitudinal section of a *GmYSL7*-RNAi 24-day-old nodule. Scale bar, 200 μ m. H. Magnification of (G) showing undeveloped (stage II) infected cells. Scale bar, 10 μ m. I. Electron microscopy of infected cells of a *GmYSL7*-RNAi 24-day-old nodule containing undeveloped single-bacteroid symbiosomes. Scale bar, 2 μ m. Yellow arrowheads indicate vacuoles. Orange arrowheads indicate symbiosomes.

Figure 4. RNAseq analysis of *GmYSL7*-RNAi nodules. A. Principal component analysis of RNAseq samples. B. Heatmap of clustering of differentially regulated genes in each nodule sample. Gene-expression values are normalized by using a z-score transformation on TPM. E1-5 indicate the RNA libraries derived from nodules transformed with empty vector and Y1 -5 indicate those transformed with the *GmYSL7* RNAi construct.

Figure 5. Gene Ontology enrichment analysis of biological processes in up and down regulated genes from *GmYSL7*-RNAi nodules. GO term enrichment analyses were performed using the ClueGO v2.5.5 plugin (Bindea et al., 2009) in Cytoscape v3.5.1 (Shannon et al., 2003). Circles represent an enriched group of genes based on their GO terms. Circle size and colour indicate the number of mapped genes and associated Term p-value corrected with Bonferroni step down.

Figure 6. *GmYSL7* does not transport iron or Fe(II)-nicotianamine (NA). DEY1530 yeast (*fet3/fet4/ftr1*) was transformed with the empty vector plasmid pDR196, *AtYSL7*, *GmYSL7* or *ZmYSL1* in pDR196GW. Serial dilutions of each yeast

transformant were applied to SD plates (that include 1.6 μM FeCl_3) with 10 μM Fe-citrate, Fe(II)-NA or no added iron (no iron) and the plates grown for 3-5 days.

Figure 7. GmYSL7 and AtYSL7 transport oligopeptides. *AtOPT4*, *AtYSL7*, *GmYSL7* in pDR196GW and the empty vector (pDR196) were introduced into the yeast *opt1* mutant, Y11213. Serial dilutions of each transformant were grown as above on minimal medium containing either 10 mM NH_4Cl (positive control) or 100 μM peptide (with sequence as indicated) as the sole source of nitrogen

Figure 8. GmYSL7 transports Syringolin A. BY4742 yeast *$\Delta\text{pdr5}::\text{KanMX6}$* transformed with the empty vector (pDR196) or the vector expressing *AtYSL7* or *GmYSL7* were plated as a lawn on solid synthetic defined (SD) media. Filter disks with the indicated SylA solutions were placed onto the plates and inhibition of growth examined after 2 days.

Figure 9. GmYSL7 is functionally equivalent to MtYSL7. A. Growth of representative wild type (WT), *ysl7-1*, and *ysl7-1* transformed with *GmYSL7* controlled by the *MtYSL7* promoter (*ysl7-1 MtYSL7_{prom}:: GmYSL7*). Bar = 1 cm. B. Dry weight of 28 dpi WT, *ysl7-1*, and *ysl7-1 MtYSL7_{prom}:: GmYSL7* plants. Data are the mean \pm SE of 5 transformed plants. C. Nitrogenase activity of 28 dpi WT, *ysl7-1*, and *ysl7-1 MtYSL7_{prom}:: GmYSL7* plants. Acetylene reduction was measured in duplicate from two sets of three-four pooled plants. Data are the mean \pm SE. * indicates statistically significant differences in Students t-test ($p < 0.05$)

LITERATURE CITED

- Abbamondi GR, De Rosa S, Iodice C, Tommonaro G (2014) Cyclic Dipeptides Produced by Marine Sponge-Associated Bacteria as Quorum Sensing Signals. *Nat. Prod. Commun* **9**: 229-232
- Acuña G, Alvarez-Morales A, Hahn M, Hennecke H (1987) A vector for the site-directed, genomic integration of foreign DNA into soybean root-nodule bacteria. *Plant Mol. Biol.* **9**: 41-50
- Agtuca BJ, Stopka SA, Evans S, Samarah L, Liu Y, Xu D, Stacey MG, Koppelaar DW, Paša-Tolić L, Anderton CR, Vertes A, Stacey G (2020) Metabolomic

- profiling of wild-type and mutant soybean root nodules using laser-ablation electrospray ionization mass spectrometry reveals altered metabolism. *Plant J.* **103**: 1937-1958. doi: 10.1111/tpj.14815.
- Avenhaus U, Cabeza RA, Liese R, Lingner A, Dittert K, Salinas-Riester G, Pommerenke C and Schulze J (2016) Short-Term Molecular Acclimation Processes of Legume Nodules to Increased External Oxygen Concentration. *Front. Plant Sci.* **6**: 1133. doi: 10.3389/fpls.2015.01133
- Becana M, Wienkoop S, Matamoros MA (2018) Sulfur Transport and Metabolism in Legume Root Nodules. *Front. Plant Sci.* **9**:1434. doi:10.3389/fpls.2018.01434
- Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, Fridman WH, Pagès F, Trajanoski Z, Galon J (2009) ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics.* **25**: 1091-1093. doi: 10.1093/bioinformatics/btp101.
- Boisson-Dernier A, Chabaud M, Garcia F, Becard G, Rosenberg C, Barker DG (2001) *Agrobacterium rhizogenes*-transformed roots of *Medicago truncatula* for the study of nitrogen-fixing and endomycorrhizal symbiotic associations. *Mol. Plant-Microbe Interact.* **14**: 695-700
- Brear EM, Day DA, Smith PMC (2013) Iron: an essential micronutrient for the legume–rhizobium symbiosis. *Front. Plant Sci.* **4**: 359
- Brear EM, Bedon F, Gavrin A, Kryvoruchko IS, Torres-Jerez I, Udvardi MK, Day DA, Smith PMC. (2020) GmVTL1a is an iron transporter on the symbiosome membrane of soybean with an important role in nitrogen fixation. *New Phytol.* **228**: 667-681. doi: 10.1111/nph.16734.
- Broughton WJ, Dilworth MJ (1971) Control of leghemoglobin synthesis in snake beans. *Biochem. J.* **125**: 1075-1080
- Cabeza R, Koester B, Liese R, Lingner A, Baumgarten V, Dirks J, Salinas-Riester G, Pommerenke C, Dittert K, Schulze J (2014) An RNA sequencing transcriptome analysis reveals novel insights into molecular aspects of the nitrate impact on the nodule activity of *Medicago truncatula*. *Plant Physiol.* **164**: 400-11. doi: 10.1104/pp.113.228312.
- Castro-Rodríguez R, Reguera M, Escudero V, Gil-Díez P, Quintana J, Prieto RI, Kumar RK, Brear EM, Grillet L, Wen J, Mysore KS, Walker EL, Smith PMC, Imperial J, González-Guerrero M (2020) *Medicago truncatula* Yellow

- Stripe-Like7* encodes a peptide transporter required for symbiotic nitrogen fixation. BioRxiv doi: <https://doi.org/10.1101/2020.03.26.009159>
- Catalano CM, Lane WS, Sherrier DJ (2004) Biochemical characterization of symbiosome membrane proteins from *Medicago truncatula* root nodules. Electrophoresis **25**: 519-531
- Chu HH, Chiecko J, Punshon T, Lanzirotti A, Lahner B, Salt DE, Walker EL (2010) Successful Reproduction Requires the Function of Arabidopsis YELLOW STRIPE-LIKE1 and YELLOW STRIPE-LIKE3 Metal-Nicotianamine Transporters in Both Vegetative and Reproductive Structures. Plant Physiol. **154**: 197-210
- Clarke VC, Loughlin PC, Day DA, Smith PMC (2014) Transport processes of the legume symbiosome membrane. Front. in Plant Sci. **5**: 699. doi:10.3389/fpls.2014.00699
- Clarke VC, Loughlin PC, Gavrin A, Chen C, Brear EM, Day DA, Smith PMC (2015) Proteomic analysis of the soybean symbiosome identifies new symbiotic proteins. Mol. Cell Proteomics **14**: 1301-1322
- Conte SS, Chu HH, Chan-Rodriguez D, Punshon T, Vasques KA, Salt DE, Walker EL (2013) *Arabidopsis thaliana* Yellow Stripe1-Like4 and Yellow Stripe1-Like6 localize to internal cellular membranes and are involved in metal ion homeostasis. Front. Plant Sci. **4**: 283. doi:10.3389/fpls.2013.00283
- Curie C, Panaviene Z, Loulergue C, Dellaporta SL, Briat JF, Walker EL (2001) Maize *yellow stripe1* encodes a membrane protein directly involved in Fe(III) uptake. Nature. **409**: 346-349. doi: 10.1038/35053080.
- Curie C, Cassin G, Couch D, Divol F, Higuchi K, Jean M, Misson J, Schikora A, Czernic P, Mari S (2009) Metal movement within the plant: contribution of nicotianamine and yellow stripe 1-like transporters. Ann. Bot. **103**: 1-11
- Dai J, Wang NQ, Xiong HC, Qiu W, Nakanishi H, Kobayashi T, Nishizawa NK, Zuo YM (2018) The Yellow Stripe-Like (YSL) Gene Functions in Internal Copper Transport in Peanut. Genes **9**: 635. doi:10.3390/genes9120635
- Divol F, Couch D, Conéjéro G, Roschzttardtz H, Mari S, Curie C (2013) The Arabidopsis YELLOW STRIPE LIKE4 and 6 Transporters Control Iron Release from the Chloroplast. Plant Cell **25**: 1040-1055

- Dohmen RJ, Strasser AWM, Honer CB, Hollenberg CP (1991) An efficient transformation procedure enabling long-term storage of competent cells of various yeast genera. *Yeast* **7**: 691-692
- Fowler D, Coyle M, Skiba U, Sutton MA, Cape JN, Reis S, Sheppard LJ, Jenkins A, Grizzetti B, Galloway JN, Vitousek P, Leach A, Bouwman AF, Butterbach-Bahl K, Dentener F, Stevenson D, Amann M, Voss M (2013) The global nitrogen cycle in the twenty-first century. *Philos. Trans. R. Soc. B-Biol. Sci.* **368**
- Gavrin A, Chiasson D, Ovchinnikova E, Kaiser BN, Bisseling T, Fedorova EE (2016) VAMP721a and VAMP721d are important for pectin dynamics and release of bacteria in soybean nodules. *New Phytol.* **210**:1011-21
- Gavrin A, Kaiser BN, Geiger D, Tyerman SD, Wen Z, Bisseling T, Fedorova EE (2014) Adjustment of Host Cells for Accommodation of Symbiotic Bacteria: Vacuole Defunctionalization, HOPS Suppression, and TIP1g Retargeting in *Medicago*. *Plant Cell* **26**: 3809-3822
- González-Guerrero M, V. E, Sáez Á, Tejada-Jiménez M (2016) Transition metal transport in plants and associated endosymbionts. Arbuscular mycorrhizal fungi and rhizobia. *Front. Plant Sci.* **7**: 1088
- Guerra D, Chapiro SM, Pratelli R, Yu S, Jia W, Leary J, Pilot G, Callis J (2017) Control of Amino Acid Homeostasis by a Ubiquitin Ligase-Coactivator Protein Complex. *J Biol Chem.* **292**: 3827-3840. doi: 10.1074/jbc.M116.766469.
- Hastwell AH, Gresshoff PM, Ferguson BJ. (2015) Genome-wide annotation and characterization of CLAVATA/ESR (CLE) peptide hormones of soybean (*Glycine max*) and common bean (*Phaseolus vulgaris*), and their orthologues of Arabidopsis thaliana. *J Exp Bot.* **66**: 5271-87. doi: 10.1093/jxb/erv351.
- Hofstetter SS, Dudnik A, Widmer H, Dudler R (2013) Arabidopsis YELLOW STRIPE-LIKE7 (YSL7) and YSL8 Transporters Mediate Uptake of Pseudomonas Virulence Factor Syringolin A into Plant Cells. *Mol. Plant-Microbe Interact.* **26**: 1302-1311
- Hohnjec N, Lenz F, Fehlberg V, Vieweg MF, Baier MC, Hause B, Küster H (2008) The Signal Peptide of the *Medicago truncatula* Modular Nodulin MtNOD25 Operates as an Address Label for the Specific Targeting of Proteins to Nitrogen-Fixing Symbiosomes. *Mol. Plant-Microbe Interact.* **22**: 63-72

- Karimi M, Inze D, Depicker A (2002) GATEWAY(TM) vectors for Agrobacterium-mediated plant transformation. *Trends Plant Sci.* **7**: 193-195
- Kiers ET, Rousseau RA, West SA, Denison RF (2003) Host sanctions and the legume-rhizobium mutualism. *Nature* **425**: 78-81
- Krusell L, Krause K, Ott T, Desbrosses G, Kramer U, Sato S, Nakamura Y, Tabata S, James EK, Sandal N, Stougaard J, Kawaguchi M, Miyamoto A, Suganuma N, Udvardi MK (2005) The sulfate transporter SST1 is crucial for symbiotic nitrogen fixation in *Lotus japonicus* root nodules. *Plant Cell* **17**: 1625-1636
- Libault M, Thibivilliers S, Bilgin DD, Radwan O, Benitez M, Clough SJ, Stacey G (2008) Identification of Four Soybean Reference Genes for Gene Expression Normalization. *Plant Genome* **1**: 44-54
- Limpens E, Ivanov S, van Esse W, Voets G, Fedorova E, Bisseling T (2009) *Medicago* N₂-Fixing Symbiosomes Acquire the Endocytic Identity Marker Rab7 but Delay the Acquisition of Vacuolar Identity. *Plant Cell* **21**: 2811-2828
- Liu T, Zeng JQ, Xia KF, Fan T, Li YG, Wang YQ, Xu XL, Zhang MY (2012) Evolutionary expansion and functional diversification of oligopeptide transporter gene family in rice. *Rice* **5**: 12. doi:10.1186/1939-8433-5-12
- Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**:550.
- Lubkowitz M (2011) The Oligopeptide Transporters: A Small Gene Family with a Diverse Group of Substrates and Functions? *Mol Plant* **4**: 407-415
- Ma C, Guo J, Kang Y, Doman K, Bryan AC, Tax FE, Yamaguchi Y, Qi Z (2014) AtPEPTIDE RECEPTOR2 mediates the AtPEPTIDE1-induced cytosolic Ca²⁺ rise, which is required for the suppression of Glutamine Dumper gene expression in Arabidopsis roots. *J Integr Plant Biol.* **56**: 684-94. doi: 10.1111/jipb.12171.
- Mohammadi-Dehcheshmeh M, Ebrahimie E, Tyerman SD, Kaiser BN (2014) A novel method based on combination of semi-in vitro and in vivo conditions in Agrobacterium rhizogenes-mediated hairy root transformation of Glycine species. *In Vitro Cell. Dev.* **50**: 282-291.
- Mohd-Noor SN, Day DA and Smith PMC (2015) Chapter 68: The Symbiosome Membrane. In: *Biological Nitrogen Fixation. Volume 2.* Ed. Frans de Bruijn. P683-694 Wiley-Blackwell. ISBN: 978-1-118-63704-3.

- Osawa H, Stacey G, Gassmann W (2006) ScOPT1 and AtOPT4 function as proton-coupled oligopeptide transporters with broad but distinct substrate specificities. *Biochem J.* **393**: 267-75. doi: 10.1042/BJ20050920.
- Patriarca EJ, Tatè R and Iaccarino M. (2002) Key role of bacterial NH₄⁺ metabolism in Rhizobium-plant symbiosis. *Microbiol Mol Biol Rev.* **66**: 203-22. doi: 10.1128/mmbr.66.2.203-222.2002. PMID: 12040124; PMCID: PMC120787.
- Pauwels L, Inze D and Goossens A (2009) Jasmonate-inducible gene: what does it mean? *Trends Plant Sci.* **14**: 87–91.
- Pessi G, Ahrens CH, Rehrauer H, Lindemann A, Hauser F, Fischer HM, Hennecke H (2007) Genome-wide transcript analysis of *Bradyrhizobium japonicum* bacteroids in soybean root nodules. *Mol. Plant-Microbe Interact.* **20**: 1353-1363
- Prell J, White JP, Bourdes A, Bunnewell S, Bongaerts RJ, Poole PS (2009) Legumes regulate Rhizobium bacteroid development and persistence by the supply of branched-chain amino acids. *Proc Natl Acad Sci U S A.* **106**: 12477-82. doi: 10.1073/pnas.0903653106.
- Prell J, Bourdès A, Kumar S, Ludwig E, Hosie A, Kinghorn S, White J, Poole P. (2010) Role of symbiotic auxotrophy in the Rhizobium-legume symbioses. *PLoS One.* **5**: e13933. doi: 10.1371/journal.pone.0013933.
- Ramakers C, Ruijter JM, Deprez RHL, Moorman AFM (2003) Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci. Lett.* **339**: 62-66
- Reid DE, Ferguson BJ, Gresshoff PM (2011) Inoculation- and nitrate-induced CLE peptides of soybean control NARK-dependent nodule formation. *Mol Plant Microbe Interact.* **24**:606-18. doi: 10.1094/MPMI-09-10-0207.
- Rentsch D, Laloi M, Rouhara I, Schmelzer E, Delrot S, Frommer WB. 1995 NTR1 encodes a high affinity oligopeptide transporter in Arabidopsis. *FEBS Lett.* **370**: 264-8.
- Saier MH (2000) Families of transmembrane transporters selective for amino acids and their derivatives. *Microbiology* **146**: 1775-1795
- Sasaki A, Yamaji N, Xia JX, Ma JF (2011) OsYSL6 Is Involved in the Detoxification of Excess Manganese in Rice. *Plant Physiol.* **157**: 1832-1840

- Schaaf G, Ludewig U, Erenoglu BE, Mori S, Kitahara T, von Wiren N (2004) ZmYS1 functions as a proton-coupled symporter for phytosiderophore- and nicotianamine-chelated metals. *J. Biol. Chem.* **279**: 9091-9096
- Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, Hyten DL, Song Q, Thelen JJ, Cheng J, Xu D, Hellsten U, May GD, Yu Y, Sakurai T, Umezawa T, Bhattacharyya MK, Sandhu D, Valliyodan B, Lindquist E, Peto M, Grant D, Shu S, Goodstein D, Barry K, Futrell-Griggs M, Abernathy B, Du J, Tian Z, Zhu L, Gill N, Joshi T, Libault M, Sethuraman A, Zhang X-C, Shinozaki K, Nguyen HT, Wing RA, Cregan P, Specht J, Grimwood J, Rokhsar D, Stacey G, Shoemaker RC, Jackson SA (2010) Genome sequence of the palaeopolyploid soybean. *Nature* **463**: 178-183
- Schneider S, Schintlmeister A, Becana M, Wagner M, Wobken D, Wienkoop S (2019) Sulfate is transported at significant rates through the symbiosome membrane and is crucial for nitrogenase biosynthesis. *Plant Cell Environ.* **42**: 1180–1189. doi:10.1111/pce.13481
- Severin AJ, Woody JL, Bolon YT, Joseph B, Diers BW, Farmer AD, Muehlbauer GJ, Nelson RT, Grant D, Specht JE, Graham MA, Cannon SB, May GD, Vance CP, Shoemaker RC (2010) RNA-Seq Atlas of *Glycine max*: A guide to the soybean transcriptome. *BMC Plant Biol.* **10** **10**: 160. doi:10.1186/1471-2229-10-160
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T (2003) Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Res* **13**: 2498–2504
- Smith PM, Atkins CA (2002) Purine biosynthesis. Big in cell division, even bigger in nitrogen assimilation. *Plant Physiol.*;128: 793–802. doi:10.1104/pp.010912
- Spizzo T, Byersdorfer C, Duesterhoeft S, Eide D. (1997) The yeast FET5 gene encodes a FET3-related multicopper oxidase implicated in iron transport. *Mol Gen Genet.* **256**: 547–556. doi:10.1007/pl00008615
- Stacey MG, Patel A, McClain WE, Mathieu M, Remley M, Rogers EE, Gassmann W, Blevins DG, Stacey G (2008) The Arabidopsis AtOPT3 protein functions in metal homeostasis and movement of iron to developing seeds. *Plant Physiol.* **146**: 589-601

- Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**: 4673–4680. doi:10.1093/nar/22.22.4673
- Trevaskis B, Wandrey M, Colebatch G, Udvardi MK (2002) The soybean *GmN6L* gene encodes a late nodulin expressed in the infected zone of nitrogen-fixing nodules. *Mol. Plant-Microbe Interact.* **15**: 630-636
- Udvardi M, Poole PS (2013) Transport and Metabolism in Legume-Rhizobia Symbioses. *Annu. Rev. Plant Biol.* **64**: 781-805
- Udvardi MK, Day DA (1997) Metabolite transport across symbiotic membranes of legume nodules. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **48**: 493-523
- Unkovich M, Baldock J (2008) Measurement of a symbiotic N₂ fixation in Australian agriculture. *Soil Biol. Biochem.* **40**: 2915-2921
- Vance CP (2001) Symbiotic nitrogen fixation and phosphorus acquisition. Plant nutrition in a world of declining renewable resources. *Plant Physiol.*, **127**: 390–397.
- Van de Velde W, Zehirov G, Szatmari A, Debreczeny M, Ishihara H, Kevei Z, Farkas A, Mikulass K, Nagy A, Tiricz H, Satiat-Jeunemaitre B, Alunni B, Bourge M, Kucho K-i, Abe M, Kereszt A, Maroti G, Uchiumi T, Kondorosi E, Mergaert P (2010) Plant Peptides Govern Terminal Differentiation of Bacteria in Symbiosis. *Science* **327**: 1122-1126
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* **3**
- Vida TA, Emr SD (1995) A new vital stain for visualizing vacuolar membrane dynamics and endocytosis in yeast. *J. Cell Biol.* **128**: 779-792
- Wienkoop S, Saalbach G (2003) Proteome analysis. Novel proteins identified at the peribacteroid membrane from *Lotus japonicus* root nodules. *Plant Physiol.* **131**: 1080-1090
- Xu C, Nadon BD, Kim KD, Jackson SA (2018) Genetic and epigenetic divergence of duplicate genes in two legume species. *Plant, Cell Environ.* **41**: 2033-2044
- Yen MR, Tseng YH, Saier MH (2001) Maize Yellow Stripe1, an iron-phytosiderophore uptake transporter, is a member of the oligopeptide transporter (OPT) family. *Microbiology* **147**: 2881-2883




Figure 1. GmYSL7 is expressed in infected cells of soybean root nodules. A. Transcript level of GmYSL7 in tissue samples from different organs. DNR, denodulated roots; nd, not detected. B. Transcript level of GmYSL7 during nodule development. 6R, roots 6 days after inoculation (DAI); 10RN, roots and nodules 10 DAI; 13N-26N, nodules the indicated DAI. Data shown are from three independent time courses. Error bars are SE ($n = 3$). Nitrogenase activity was first detected at 18 DAI. The relative expression was quantified by RT-qPCR and normalized to a soybean ubiquitin gene (GmUBI3, Glyma20g27950.1). C. Transgenic root expressing pYSL7:GFP-GUS. GUS staining was not detectable in the very early stages of nodule development. Arrowhead indicates a nodule initiation. D. Transgenic pYSL7:GFP-GUS 10-day-old nodule primordia. IR, infected region. E. Transgenic pYSL7:GFP-GUS mature nodule. GUS staining is restricted to infected cells. OC, outer cortex. F. Magnification of E. Scale bars, 150 μ m.




Figure 2. Localization of GmYSL7 in soybean nodule cells infected with rhizobia. A. Localization of GFP-GmYSL7 on symbiosome membranes in an infected cell counterstained by membrane lipophilic dye FM4-64; IC, infected cell; NI, non-infected cell; ROI, region of interest. B. Fluorescent intensity plot of ROI from A. C. Colocalization analysis of GFP-YSL7 with FM4-64. The colocalised points are highlighted by white in colocalization map. D. Scatterplot between the intensities of the GFP-YSL7 and FM4-64 in C. E. Superimposed confocal z-stack sections of GFP-GmYSL7 signal in rhizobia-infected nodule cells. F. Free GFP localizes to the cytoplasmic spaces surrounding symbiosomes in infected cells. G. MtNOD25-GFP (Hohnjec et al. 2009) localizes to the peribacteroid space inside the symbiosomes. Scale bars, 5 μ m.




Figure 3. RNAi silencing of *GmYSL7* affects nodule development. A. Transcript level of *GmYSL7* and its closest homologs in 24-day-old nodules of empty vector control and *GmYSL7*-RNAi plants (error bars represent SD; n=4; t-test: *, p<0.05). B. Nitrogenase enzyme activity in 24-day-old nodules of empty vector control and *GmYSL7*-RNAi plants (error bars represent SD; n=8; t-test: **, p<0.01). C. Fresh weight of 24-day-old nodules of empty vector control and RNAi-*GmYSL7* (error bars represent SD; t-test: *, p<0.05). D. Longitudinal section of a 24-day-old nodule from an empty vector control plant. Scale bar, 200 μ m. E. Magnification of (D) showing developed (stage IV) infected cells. Scale bar, 10 μ m. F. Electron microscopy of infected cells of a 24-day-old nodule from empty vector control containing developed multibacteroid symbiosomes. Scale bar, 2 μ m. G. Longitudinal section of a *GmYSL7*-RNAi 24-day-old nodule. Scale bar, 200 μ m. H. Magnification of (G) showing undeveloped (stage II) infected cells. Scale bar, 10 μ m. I. Electron microscopy of infected cells of a *GmYSL7*-RNAi 24-day-old nodule containing undeveloped single-bacteroid symbiosomes. Scale bar, 2 μ m. Yellow arrowheads indicate vacuoles. Orange arrowheads indicate symbiosomes.




Figure 4. RNaseq analysis of *GmYSL7*-RNAi nodules. A. Principal component analysis of RNaseq samples. B. Heatmap of clustering of differentially regulated genes in each nodule sample. E1-5 indicate the RNA libraries derived from nodules transformed with empty vector and Y1 -5 indicate those transformed with the *GmYSL7* RNAi construct.




Figure 5. Gene Ontology enrichment analysis of biological processes in up (A) and down (B) regulated genes from *GmYSL7*-RNAi nodules. GO term enrichment analyses were performed using the ClueGO v2.5.5 plugin (Bindea et al., 2009) in Cytoscape v3.5.1 (Shannon et al., 2003). Circles represent an enriched group of genes based on their GO terms. Circle size and colour indicate the number of mapped genes and associated Term p-value corrected with Bonferroni step down.




Figure 6. GmYSL7 does not transport Fe(II)-nicotianamine (NA). DEY1530 yeast (*fet3/fet4/ftr1*) was transformed with the empty vector plasmid pDR196, *AtYSL7*, *GmYSL7*, *ZmYS1* in pDR196-GW. Serial dilutions of each yeast transformant were applied to SD plates (that include 1.6 μM FeCl_3) with 10 μM Fe-citrate, Fe(II)-NA or no added iron (no iron) and the plates grown for 3-5 days.




Figure 7. GmYSL7 and AtYSL7 transport oligopeptides. *AtOPT4*, *AtYSL7*, *GmYSL7* in pDR196-GW and the empty vector (pDR196) were introduced into the yeast *opt1* mutant, Y11213. Serial dilutions of each transformant were grown as above on minimal medium containing either 10 mM NH_4Cl (positive control) or 100 μM peptide (with sequence as indicated) as the sole source of nitrogen.




Figure 8. GmYSL7 transports Syringolin A. BY4742 yeast $\Delta pr5:KanMX6$ transformed with the empty vector (pDR196) or the vector expressing *AtYSL7* or *GmYSL7* were plated as a lawn on solid synthetic defined (SD) media. Filter disks with the indicated SylA solutions were placed onto the plates and inhibition of growth examined after 2 days.



A**B****C**

Figure 9. GmYSL7 is functionally equivalent to MtYSL7. A. Growth of representative wild type (WT), *ysl7-1*, and *ysl7-1* transformed with *GmYSL7* controlled by the *MtYSL7* promoter (*ysl7-1* *MtYSL7_{prom}::GmYSL7*). Bar = 1 cm. B. Dry weight of 28 dpi WT, *ysl7-1*, and *ysl7-1* *MtYSL7_{prom}::GmYSL7* plants. Data are the mean \pm SE of 5 transformed plants. C. Nitrogenase activity of 28 dpi WT, *ysl7-1*, and *ysl7-1* *MtYSL7_{prom}::GmYSL7* plants. Acetylene reduction was measured in duplicate from two sets of three-four pooled plants. Data are the mean \pm SE. * indicates statistically significant differences in Students t-test ($p < 0.05$)

Parsed Citations

Abbamondi GR, De Rosa S, Iodice C, Tommonaro G (2014) Cyclic Dipeptides Produced by Marine Sponge-Associated Bacteria as Quorum Sensing Signals. *Nat. Prod. Commun* 9: 229-232

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Acuña G, Alvarez-Morales A, Hahn M, Hennecke H (1987) A vector for the site-directed, genomic integration of foreign DNA into soybean root-nodule bacteria. *Plant Mol. Biol.* 9: 41-50

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Agtuca BJ, Stopka SA, Evans S, Samarah L, Liu Y, Xu D, Stacey MG, Koppelaar DW, Paša-Tolić L, Anderton CR, Vertes A, Stacey G (2020) Metabolomic profiling of wild-type and mutant soybean root nodules using laser-ablation electrospray ionization mass spectrometry reveals altered metabolism. *Plant J.* 103: 1937-1958. doi: 10.1111/tj.14815.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Avenhaus U, Cabeza RA, Liese R, Lingner A, Dittert K, Salinas-Riester G, Pommerenke C and Schulze J (2016) Short-Term Molecular Acclimation Processes of Legume Nodules to Increased External Oxygen Concentration. *Front. Plant Sci.* 6: 1133. doi: 10.3389/fpls.2015.01133

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Becana M, Wienkoop S, Matamoros MA (2018) Sulfur Transport and Metabolism in Legume Root Nodules. *Front. Plant Sci.* 9 :1434. doi:10.3389/fpls.2018.01434

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, Fridman WH, Pagès F, Trajanoski Z, Galon J (2009) ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics.* 25: 1091-1093. doi: 10.1093/bioinformatics/btp101.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Boisson-Dernier A, Chabaud M, Garcia F, Becard G, Rosenberg C, Barker DG (2001) Agrobacterium rhizogenes-transformed roots of *Medicago truncatula* for the study of nitrogen-fixing and endomycorrhizal symbiotic associations. *Mol. Plant-Microbe Interact.* 14: 695-700

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Brear EM, Day DA, Smith PMC (2013) Iron: an essential micronutrient for the legume–rhizobium symbiosis. *Front. Plant Sci.* 4: 359

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Brear EM, Bedon F, Gavrin A, Kryvoruchko IS, Torres-Jerez I, Udvardi MK, Day DA, Smith PMC. (2020) GmVTL1a is an iron transporter on the symbiosome membrane of soybean with an important role in nitrogen fixation. *New Phytol.* 228: 667-681. doi: 10.1111/nph.16734.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Broughton WJ, Dilworth MJ (1971) Control of leghemoglobin synthesis in snake beans. *Biochem. J.* 125: 1075-1080

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cabeza R, Koester B, Liese R, Lingner A, Baumgarten V, Dirks J, Salinas-Riester G, Pommerenke C, Dittert K, Schulze J (2014) An RNA sequencing transcriptome analysis reveals novel insights into molecular aspects of the nitrate impact on the nodule activity of *Medicago truncatula*. *Plant Physiol.* 164: 400-11. doi: 10.1104/pp.113.228312.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Castro-Rodríguez R, Reguera M, Escudero V, Gil-Díez P, Quintana J, Prieto RI, Kumar RK, Brear EM, Grillet L, Wen J, Mysore KS, Walker EL, Smith PMC, Imperial J, González-Guerrero M (2020) *Medicago truncatula* Yellow Stripe-Like7 encodes a peptide transporter required for symbiotic nitrogen fixation. *BioRxiv* doi: <https://doi.org/10.1101/2020.03.26.009159>

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Catalano CM, Lane WS, Sherrier DJ (2004) Biochemical characterization of symbiosome membrane proteins from *Medicago truncatula* root nodules. *Electrophoresis* 25: 519-531

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chu HH, Chiecko J, Punshon T, Lanzirotti A, Lahner B, Salt DE, Walker EL (2010) Successful Reproduction Requires the Function of Arabidopsis YELLOW STRIPE-LIKE1 and YELLOW STRIPE-LIKE3 Metal-Nicotianamine Transporters in Both Vegetative and Reproductive Structures. *Plant Physiol.* 154: 197-210

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Clarke VC, Loughlin PC, Day DA, Smith PMC (2014) Transport processes of the legume symbiosome membrane. *Front. in Plant Sci.* 5: 699. doi:10.3389/fpls.2014.00699

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Clarke VC, Loughlin PC, Gavrin A, Chen C, Brear EM, Day DA, Smith PMC (2015) Proteomic analysis of the soybean symbiosome identifies new symbiotic proteins. *Mol. Cell Proteomics* 14: 1301-1322

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Conte SS, Chu HH, Chan-Rodriguez D, Punshon T, Vasques KA, Salt DE, Walker EL (2013) Arabidopsis thaliana Yellow Stripe1-Like4

and Yellow Stripe1-Like6 localize to internal cellular membranes and are involved in metal ion homeostasis. *Front. Plant Sci.* 4: 283. doi:10.3389/fpls.2013.00283

Google Scholar: [Author Only Title Only Author and Title](#)

Curie C, Panaviene Z, Loulergue C, Dellaporta SL, Briat JF, Walker EL (2001) Maize yellow stripe1 encodes a membrane protein directly involved in Fe(III) uptake. *Nature*. 409: 346-349. doi: 10.1038/35053080.

Google Scholar: [Author Only Title Only Author and Title](#)

Curie C, Cassin G, Couch D, Divol F, Higuchi K, Jean M, Misson J, Schikora A, Czernic P, Mari S (2009) Metal movement within the plant: contribution of nicotianamine and yellow stripe 1-like transporters. *Ann. Bot.* 103: 1-11

Google Scholar: [Author Only Title Only Author and Title](#)

Dai J, Wang NQ, Xiong HC, Qiu W, Nakanishi H, Kobayashi T, Nishizawa NK, Zuo YM (2018) The Yellow Stripe-Like (YSL) Gene Functions in Internal Copper Transport in Peanut. *Genes* 9: 635. doi:10.3390/genes9120635

Google Scholar: [Author Only Title Only Author and Title](#)

Divol F, Couch D, Conéjéro G, Roschztardt H, Mari S, Curie C (2013) The Arabidopsis YELLOW STRIPE LIKE4 and 6 Transporters Control Iron Release from the Chloroplast. *Plant Cell* 25: 1040-1055

Google Scholar: [Author Only Title Only Author and Title](#)

Dohmen RJ, Strasser AWM, Honer CB, Hollenberg CP (1991) An efficient transformation procedure enabling long-term storage of competent cells of various yeast genera. *Yeast* 7: 691-692

Google Scholar: [Author Only Title Only Author and Title](#)

Fowler D, Coyle M, Skiba U, Sutton MA, Cape JN, Reis S, Sheppard LJ, Jenkins A, Grizzetti B, Galloway JN, Vitousek P, Leach A, Bouwman AF, Butterbach-Bahl K, Dentener F, Stevenson D, Amann M, Voss M (2013) The global nitrogen cycle in the twenty-first century. *Philos. Trans. R. Soc. B-Biol. Sci.* 368

Google Scholar: [Author Only Title Only Author and Title](#)

Gavrin A, Chiasson D, Ovchinnikova E, Kaiser BN, Bisseling T, Fedorova EE (2016) VAMP721a and VAMP721d are important for pectin dynamics and release of bacteria in soybean nodules. *New Phytol.* 210:1011-21

Google Scholar: [Author Only Title Only Author and Title](#)

Gavrin A, Kaiser BN, Geiger D, Tyerman SD, Wen Z, Bisseling T, Fedorova EE (2014) Adjustment of Host Cells for Accommodation of Symbiotic Bacteria: Vacuole Defunctionalization, HOPS Suppression, and TIP1g Retargeting in Medicago. *Plant Cell* 26: 3809-3822

Google Scholar: [Author Only Title Only Author and Title](#)

González-Guerrero M, V. E, Sáez Á, Tejada-Jiménez M (2016) Transition metal transport in plants and associated endosymbionts. *Arbuscular mycorrhizal fungi and rhizobia*. *Front. Plant Sci.* 7: 1088

Google Scholar: [Author Only Title Only Author and Title](#)

Guerra D, Chapiro SM, Pratelli R, Yu S, Jia W, Leary J, Pilot G, Callis J (2017) Control of Amino Acid Homeostasis by a Ubiquitin Ligase-Coactivator Protein Complex. *J Biol Chem*. 292: 3827-3840. doi: 10.1074/jbc.M116.766469.

Google Scholar: [Author Only Title Only Author and Title](#)

Hastwell AH, Gresshoff PM, Ferguson BJ. (2015) Genome-wide annotation and characterization of CLAVATA/ESR (CLE) peptide hormones of soybean (*Glycine max*) and common bean (*Phaseolus vulgaris*), and their orthologues of *Arabidopsis thaliana*. *J Exp Bot.* 66: 5271-87. doi: 10.1093/jxb/erv351.

Google Scholar: [Author Only Title Only Author and Title](#)

Hofstetter SS, Dudnik A, Widmer H, Dudler R (2013) Arabidopsis YELLOW STRIPE-LIKE7 (YSL7) and YSL8 Transporters Mediate Uptake of Pseudomonas Virulence Factor Syringolin A into Plant Cells. *Mol. Plant-Microbe Interact.* 26: 1302-1311

Google Scholar: [Author Only Title Only Author and Title](#)

Hohnjec N, Lenz F, Fehlberg V, Vieweg MF, Baier MC, Hause B, Küster H (2008) The Signal Peptide of the Medicago truncatula Modular Nodulin MtNOD25 Operates as an Address Label for the Specific Targeting of Proteins to Nitrogen-Fixing Symbiosomes. *Mol. Plant-Microbe Interact.* 22: 63-72

Google Scholar: [Author Only Title Only Author and Title](#)

Karimi M, Inze D, Depicker A (2002) GATEWAY(TM) vectors for Agrobacterium-mediated plant transformation. *Trends Plant Sci.* 7: 193-195

Google Scholar: [Author Only Title Only Author and Title](#)

Kiers ET, Rousseau RA, West SA, Denison RF (2003) Host sanctions and the legume-rhizobium mutualism. *Nature* 425: 78-81

Google Scholar: [Author Only Title Only Author and Title](#)

Krusell L, Krause K, Ott T, Desbrosses G, Kramer U, Sato S, Nakamura Y, Tabata S, James EK, Sandal N, Stougaard J, Kawaguchi M, Miyamoto A, Sukanuma N, Udvardi MK (2005) The sulfate transporter SST1 is crucial for symbiotic nitrogen fixation in Lotus japonicus root nodules. *Plant Cell* 17: 1625-1636

Google Scholar: [Author Only Title Only Author and Title](#)

Libault M, Thibivilliers S, Bilgin DD, Radwan O, Benitez M, Clough SJ, Stacey G (2008) Identification of Four Soybean Reference Genes for Gene Expression Normalization. *Plant Genome* 1: 44-54

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Limpens E, Ivanov S, van Esse W, Voets G, Fedorova E, Bisseling T (2009) Medicago N₂-Fixing Symbiosomes Acquire the Endocytic Identity Marker Rab7 but Delay the Acquisition of Vacuolar Identity. *Plant Cell* 21: 2811-2828

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu T, Zeng JQ, Xia KF, Fan T, Li YG, Wang YQ, Xu XL, Zhang MY (2012) Evolutionary expansion and functional diversification of oligopeptide transporter gene family in rice. *Rice* 5: 12. doi:10.1186/1939-8433-5-12

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15:550.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lubkowitz M (2011) The Oligopeptide Transporters: A Small Gene Family with a Diverse Group of Substrates and Functions? *Mol Plant* 4: 407-415

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ma C, Guo J, Kang Y, Doman K, Bryan AC, Tax FE, Yamaguchi Y, Qi Z (2014) AtPEPTIDE RECEPTOR2 mediates the AtPEPTIDE1-induced cytosolic Ca²⁺ rise, which is required for the suppression of Glutamine Dumper gene expression in Arabidopsis roots. *J Integr Plant Biol*. 56: 684-94. doi: 10.1111/jipb.12171.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mohammadi-Dehcheshmeh M, Ebrahimie E, Tyerman SD, Kaiser BN (2014) A novel method based on combination of semi-in vitro and in vivo conditions in Agrobacterium rhizogenes-mediated hairy root transformation of Glycine species. *In Vitro Cell. Dev.* 50: 282-291.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mohd-Noor SN, Day DA and Smith PMC (2015) Chapter 68: The Symbiosome Membrane. In: Biological Nitrogen Fixation. Volume 2. Ed. Frans de Bruijn. P683-694 Wiley-Blackwell. ISBN: 978-1-118-63704-3.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Osawa H, Stacey G, Gassmann W (2006) ScOPT1 and AtOPT4 function as proton-coupled oligopeptide transporters with broad but distinct substrate specificities. *Biochem J*. 393: 267-75. doi: 10.1042/BJ20050920.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Patriarca EJ, Tatè R and Iaccarino M. (2002) Key role of bacterial NH₄⁺ metabolism in Rhizobium-plant symbiosis. *Microbiol Mol Biol Rev*. 66: 203-22. doi: 10.1128/mmr.66.2.203-222.2002. PMID: 12040124; PMCID: PMC120787.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pauwels L, Inze D and Goossens A (2009) Jasmonate-inducible gene:

what does it mean? Trends Plant Sci. 14: 87–91.

Pessi G, Ahrens CH, Rehrauer H, Lindemann A, Hauser F, Fischer HM, Hennecke H (2007) Genome-wide transcript analysis of Bradyrhizobium japonicum bacteroids in soybean root nodules. *Mol. Plant-Microbe Interact*. 20: 1353-1363

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Prell J, White JP, Bourdes A, Bunnewell S, Bongaerts RJ, Poole PS (2009) Legumes regulate Rhizobium bacteroid development and persistence by the supply of branched-chain amino acids. *Proc Natl Acad Sci U S A*. 106: 12477-82. doi: 10.1073/pnas.0903653106.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Prell J, Bourdès A, Kumar S, Lodwig E, Hosie A, Kinghorn S, White J, Poole P. (2010) Role of symbiotic auxotrophy in the Rhizobium-legume symbioses. *PLoS One*. 5: e13933. doi: 10.1371/journal.pone.0013933.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ramakers C, Ruijter JM, Deprez RHL, Moorman AFM (2003) Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci. Lett*. 339: 62-66

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Reid DE, Ferguson BJ, Gresshoff PM (2011) Inoculation- and nitrate-induced CLE peptides of soybean control NARK-dependent nodule formation. *Mol Plant Microbe Interact*. 24:606-18. doi: 10.1094/MPMI-09-10-0207.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rentsch D, Laloi M, Rouhara I, Schmelzer E, Delrot S, Frommer WB. 1995 NTR1 encodes a high affinity oligopeptide transporter in Arabidopsis. FEBS Lett. 370: 264-8.

Saier MH (2000) Families of transmembrane transporters selective for amino acids and their derivatives. *Microbiology* 146: 1775-1795

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sasaki A, Yamaji N, Xia JX, Ma JF (2011) OsYSL6 Is Involved in the Detoxification of Excess Manganese in Rice. *Plant Physiol*. 157: 1832-1840

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schaaf G, Ludewig U, Erenoglu BE, Mori S, Kitahara T, von Wren N (2004) ZmYS1 functions as a proton-coupled symporter for

phytosiderophore- and nicotianamine-chelated metals. *J. Biol. Chem.* 279: 9091-9096

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, Hyten DL, Song Q, Thelen JJ, Cheng J, Xu D, Hellsten U, May GD, Yu Y, Sakurai T, Umezawa T, Bhattacharyya MK, Sandhu D, Valliyodan B, Lindquist E, Peto M, Grant D, Shu S, Goodstein D, Barry K, Futrell-Griggs M, Abernathy B, Du J, Tian Z, Zhu L, Gill N, Joshi T, Libault M, Sethuraman A, Zhang X-C, Shinozaki K, Nguyen HT, Wing RA, Cregan P, Specht J, Grimwood J, Rokhsar D, Stacey G, Shoemaker RC, Jackson SA (2010) Genome sequence of the palaeopolyploid soybean. *Nature* 463: 178-183

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schneider S, Schintlmeister A, Becana M, Wagner M, Wobken D, Wienkoop S (2019) Sulfate is transported at significant rates through the symbiosome membrane and is crucial for nitrogenase biosynthesis. *Plant Cell Environ.* 42: 1180–1189. doi:10.1111/pce.13481

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Severin AJ, Woody JL, Bolon YT, Joseph B, Diers BW, Farmer AD, Muehlbauer GJ, Nelson RT, Grant D, Specht JE, Graham MA, Cannon SB, May GD, Vance CP, Shoemaker RC (2010) RNA-Seq Atlas of Glycine max: A guide to the soybean transcriptome. *BMC Plant Biol.* 10 10: 160. doi:10.1186/1471-2229-10-160

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T (2003) Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Res* 13: 2498–2504

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Smith PM, Atkins CA (2002) Purine biosynthesis. Big in cell division, even bigger in nitrogen assimilation. *Plant Physiol.* 128: 793–802. doi:10.1104/pp.010912

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Spizzo T, Byersdorfer C, Dueterhoeft S, Eide D. (1997) The yeast FET5 gene encodes a FET3-related multicopper oxidase implicated in iron transport. *Mol Gen Genet.* 256: 547–556. doi:10.1007/pl00008615

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Stacey MG, Patel A, McClain WE, Mathieu M, Remley M, Rogers EE, Gassmann W, Blevins DG, Stacey G (2008) The Arabidopsis AtOPT3 protein functions in metal homeostasis and movement of iron to developing seeds. *Plant Physiol.* 146: 589-601

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22: 4673–4680. doi:10.1093/nar/22.22.4673

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Trevaskis B, Wandrey M, Colebatch G, Udvardi MK (2002) The soybean GmN6L gene encodes a late nodulin expressed in the infected zone of nitrogen-fixing nodules. *Mol. Plant-Microbe Interact.* 15: 630-636

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Udvardi M, Poole PS (2013) Transport and Metabolism in Legume-Rhizobia Symbioses. *Annu. Rev. Plant Biol.* 64: 781-805

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Udvardi MK, Day DA (1997) Metabolite transport across symbiotic membranes of legume nodules. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 48: 493-523

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Unkovich M, Baldock J (2008) Measurement of a symbiotic N₂ fixation in Australian agriculture. *Soil Biol. Biochem.* 40: 2915-2921

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vance CP (2001) Symbiotic nitrogen fixation and phosphorus acquisition. Plant nutrition in a world of declining renewable resources. *Plant Physiol.* 127: 390–397.

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Van de Velde W, Zehirov G, Szatmari A, Debreczeny M, Ishihara H, Kevei Z, Farkas A, Mikulass K, Nagy A, Tiricz H, Satiat-Jeunemaitre B, Alunni B, Bourge M, Kucho K-i, Abe M, Kereszt A, Maroti G, Uchiumi T, Kondorosi E, Mergaert P (2010) Plant Peptides Govern Terminal Differentiation of Bacteria in Symbiosis. *Science* 327: 1122-1126

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* 3

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vida TA, Emr SD (1995) A new vital stain for visualizing vacuolar membrane dynamics and endocytosis in yeast. *J. Cell Biol.* 128: 779-792

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wienkoop S, Saalbach G (2003) Proteome analysis. Novel proteins identified at the peribacteroid membrane from *Lotus japonicus* root nodules. *Plant Physiol.* 131: 1080-1090

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xu C, Nadon BD, Kim KD, Jackson SA (2018) Genetic and epigenetic divergence of duplicate genes in two legume species. *Plant, Cell Environ.* 41: 2033-2044

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yen MR, Tseng YH, Saier MH (2001) Maize Yellow Stripe1, an iron-phytosiderophore uptake transporter, is a member of the oligopeptide transporter (OPT) family. *Microbiology* 147: 2881-2883

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)